

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
16 May 2002 (16.05.2002)

PCT

(10) International Publication Number  
**WO 02/37967 A1**

(51) International Patent Classification<sup>7</sup>: **A01N 63/00**,  
A61K 39/395, 48/00, C07H 21/04, C12N 5/00, 15/00,  
C12P 21/06

(74) Agents: CALDWELL, John, W. et al.; Woodcock Wash-  
burn Kurtz MacKiewicz & Norris LLP, 46th floor, One Lib-  
erty Place, Philadelphia, PA 19103 (US).

(21) International Application Number: PCT/US00/30588

(22) International Filing Date:

7 November 2000 (07.11.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(71) Applicant (for all designated States except US): **MOR-  
PHOTEK INC.** [US/US]; Suite 516, 3624 Market Street,  
Philadelphia, PA 19104 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, MI, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **NICOLAIDES,**  
**Nicholas, C.** [US/US]; 4 Cider Mill Court, Boothwyn, PA  
19061 (US). **GRASSO, Luigi** [US/US]; Apartment #816,  
834 Chestnut Street, Philadelphia, PA 19107 (US). **SASS,**  
**Philip, M.** [US/US]; 1903 Blackhawk Circle, Audubon,  
PA 19403 (US).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: METHODS FOR GENERATING GENETICALLY ALTERED ANTIBODY-PRODUCING CELL LINES WITH IM-  
PROVED ANTIBODY CHARACTERISTICS

(57) Abstract: Dominant negative alleles of human mismatch repair genes can be used to generate hypermutable cells and or-  
ganisms. By introducing these genes into cells and transgenic animals, new cell lines and animal varieties with novel and useful  
properties can be prepared more efficiently than by relying on the natural rate of mutation. These methods are useful for generating  
genetic diversity within immunoglobulin genes directed against an antigen of interest to produce altered antibodies with enhanced  
biochemical activity. Moreover, these methods are useful for generating antibody-producing cells with increased level of antibody  
production.



WO 02/37967 A1

**METHODS FOR GENERATING GENETICALLY ALTERED ANTIBODY-  
PRODUCING CELL LINES WITH IMPROVED ANTIBODY  
CHARACTERISTICS**

**5 TECHNICAL FIELD OF THE INVENTION**

The invention is related to the area of antibody maturation and cellular production. In particular, it is related to the field of mutagenesis.

**BACKGROUND OF THE INVENTION**

10 The use of antibodies to block the activity of foreign and/or endogenous polypeptides provides an effective and selective strategy for treating the underlying cause of disease. In particular is the use of monoclonal antibodies (MAb) as effective therapeutics such as the FDA approved ReoPro (Glaser, V. (1996) Can ReoPro repolish tarnished monoclonal therapeutics? *Nat. Biotechnol.* 14:1216-1217), an anti-platelet MAb  
15 from Centocor; Herceptin (Weiner, L.M. (1999) Monoclonal antibody therapy of cancer. *Semin. Oncol.* 26:43-51), an anti-Her2/neu MAb from Genentech; and Synagis (Saez-Llorens, X.E., *et al.* (1998) Safety and pharmacokinetics of an intramuscular humanized monoclonal antibody to respiratory syncytial virus in premature infants and infants with bronchopulmonary dysplasia. *Pediat. Infect. Dis. J.* 17:787-791), an anti-respiratory  
20 syncytial virus MAb produced by Medimmune.

Standard methods for generating MAbs against candidate protein targets are known by those skilled in the art. Briefly, rodents such as mice or rats are injected with a purified antigen in the presence of adjuvant to generate an immune response (Shield, C.F., *et al.* (1996) A cost-effective analysis of OKT3 induction therapy in cadaveric kidney  
25 transplantation. *Am. J. Kidney Dis.* 27:855-864). Rodents with positive immune sera are sacrificed and splenocytes are isolated. Isolated splenocytes are fused to melanomas to produce immortalized cell lines that are then screened for antibody production. Positive lines are isolated and characterized for antibody production. The direct use of rodent MAbs as human therapeutic agents were confounded by the fact that human anti-rodent  
30 antibody (HARA) responses occurred in a significant number of patients treated with the rodent-derived antibody (Khazaeli, M.B., *et al.*, (1994) Human immune response to monoclonal antibodies. *J. Immunother.* 15:42-52). In order to circumvent the problem of HARA, the grafting of the complementarity determining regions (CDRs), which are the

critical motifs found within the heavy and light chain variable regions of the immunoglobulin (Ig) subunits making up the antigen binding domain, onto a human antibody backbone found these chimeric molecules are able to retain their binding activity to antigen while lacking the HARA response (Emery, S.C., and Harris, W.J. "Strategies  
5 for humanizing antibodies" In: ANTIBODY ENGINEERING C.A.K. Borrebaeck (Ed.) Oxford University Press, N.Y. 1995. pp. 159-183. A common problem that exists during the "humanization" of rodent-derived MAbs (referred to hereon as HAb) is the loss of binding affinity due to conformational changes in the 3 dimensional structure of the CDR domain upon grafting onto the human Ig backbone (U.S. Patent No. 5,530,101 to Queen *et al.*). To  
10 overcome this problem, additional HAb vectors are usually needed to be engineered by inserting or deleting additional amino acid residues within the framework region and/or within the CDR coding region itself in order to recreate high affinity HAbs (U.S. Patent No. 5,530,101 to Queen *et al.*). This process is a very time consuming procedure that involves the use of expensive computer modeling programs to predict changes that may  
15 lead to a high affinity HAb. In some instances the affinity of the HAb is never restored to that of the MAb, rendering them of little therapeutic use.

Another problem that exists in antibody engineering is the generation of stable, high yielding producer cell lines that is required for manufacturing of the molecule for clinical materials. Several strategies have been adopted in standard practice by those  
20 skilled in the art to circumvent this problem. One method is the use of Chinese Hamster Ovary (CHO) cells transfected with exogenous Ig fusion genes containing the grafted human light and heavy chains to produce whole antibodies or single chain antibodies, which are a chimeric molecule containing both light and heavy chains that form an antigen-binding polypeptide (Reff, M.E. (1993) High-level production of recombinant immunoglobulins in mammalian cells. *Curr. Opin. Biotechnol.* 4:573-576). Another  
25 method employs the use of human lymphocytes derived from transgenic mice containing a human grafted immune system or transgenic mice containing a human Ig gene repertoire. Yet another method employs the use of monkeys to produce primate MAbs, which have been reported to lack a human anti-monkey response (Neuberger, M., and Gruggermann, M. (1997) Monoclonal antibodies. Mice perform a human repertoire. *Nature* 386:25-26).  
30 In all cases, the generation of a cell line that is capable of generating sufficient amounts of high affinity antibody poses a major limitation for producing sufficient materials for clinical studies. Because of these limitations, the utility of other recombinant systems such

as plants are currently being explored as systems that will lead to the stable, high-level production of humanized antibodies (Fiedler, U., and Conrad, U. (1995) High-level production and long-term storage of engineered antibodies in transgenic tobacco seeds. *Bio/Technology* 13:1090-1093).

5           A method for generating diverse antibody sequences within the variable domain that results in HAbs and MAbs with high binding affinities to antigens would be useful for the creation of more potent therapeutic and diagnostic reagents respectively. Moreover, the generation of randomly altered nucleotide and polypeptide residues throughout an entire antibody molecule will result in new reagents that are less antigenic and/or have  
10   beneficial pharmacokinetic properties. The invention described herein is directed to the use of random genetic mutation throughout an antibody structure *in vivo* by blocking the endogenous mismatch repair (MMR) activity of a host cell producing immunoglobulins that encode biochemically active antibodies. The invention also relates to methods for repeated *in vivo* genetic alterations and selection for antibodies with enhanced binding and  
15   pharmacokinetic profiles.

          In addition, the ability to develop genetically altered host cells that are capable of secreting increased amounts of antibody will also provide a valuable method for creating cell hosts for product development. The invention described herein is directed to the creation of genetically altered cell hosts with increased antibody production via the  
20   blockade of MMR.

          The invention facilitates the generation of high affinity antibodies and the production of cell lines with elevated levels of antibody production. Other advantages of the present invention are described in the examples and figures described herein.

#### **SUMMARY OF THE INVENTION**

25           The invention provides methods for generating genetically altered antibodies (including single chain molecules) and antibody producing cell hosts *in vitro* and *in vivo*, whereby the antibody possess a desired biochemical property(s), such as, but not limited to, increased antigen binding, increased gene expression, and/or enhanced extracellular secretion by the cell host. One method for identifying antibodies with increased binding  
30   activity or cells with increased antibody production is through the screening of MMR defective antibody producing cell clones that produce molecules with enhanced binding properties or clones that have been genetically altered to produce enhanced amounts of antibody product.

The antibody producing cells suitable for use in the invention include, but are not limited to rodent, primate, or human hybridomas or lymphoblastoids; mammalian cells transfected and expressing exogenous Ig subunits or chimeric single chain molecules; plant cells, yeast or bacteria transfected and expressing exogenous Ig subunits or chimeric single chain molecules.

Thus, the invention provides methods for making hypermutable antibody-producing cells by introducing a polynucleotide comprising a dominant negative allele of a mismatch repair gene into cells that are capable of producing antibodies. The cells that are capable of producing antibodies include cells that naturally produce antibodies, and cells that are engineered to produce antibodies through the introduction of immunoglobulin encoding sequences. Conveniently, the introduction of polynucleotide sequences into cells is accomplished by transfection.

The invention also provides methods of making hypermutable antibody producing cells by introducing a dominant negative mismatch repair (MMR) gene such as *PMS2* (preferably human *PMS2*), *MLH1*, *PMS1*, *MSH2*, or *MSH2* into cells that are capable of producing antibodies. The dominant negative allele of a mismatch repair gene may be a truncation mutation of a mismatch repair gene (preferably a truncation mutation at codon 134, or a thymidine at nucleotide 424 of wild-type *PMS2*). The invention also provides methods in which mismatch repair gene activity is suppressed. This may be accomplished, for example, using antisense molecules directed against the mismatch repair gene or transcripts.

Other embodiments of the invention provide methods for making a hypermutable antibody producing cells by introducing a polynucleotide comprising a dominant negative allele of a mismatch repair gene into fertilized eggs of animals. These methods may also include subsequently implanting the eggs into pseudo-pregnant females whereby the fertilized eggs develop into a mature transgenic animal. The mismatch repair genes may include, for example, *PMS2* (preferably human *PMS2*), *MLH1*, *PMS1*, *MSH2*, or *MSH2*. The dominant negative allele of a mismatch repair gene may be a truncation mutation of a mismatch repair gene (preferably a truncation mutation at codon 134, or a thymidine at nucleotide 424 of wild-type *PMS2*).

The invention further provides homogeneous compositions of cultured, hypermutable, mammalian cells that are capable of producing antibodies and contain a dominant negative allele of a mismatch repair gene. The mismatch repair genes may

include, for example, *PMS2* (preferably human *PMS2*), *MLH1*, *PMS1*, *MSH2*, or *MSH2*. The dominant negative allele of a mismatch repair gene may be a truncation mutation of a mismatch repair gene (preferably a truncation mutation at codon 134, or a thymidine at nucleotide 424 of wild-type *PMS2*). The cells of the culture may contain *PMS2*, (preferably human *PMS2*), *MLH1*, or *PMS1*; or express a human *mutL* homolog, or the first 133 amino acids of hPMS2.

The invention further provides methods for generating a mutation in an immunoglobulin gene of interest by culturing an immunoglobulin producing cell selected for an immunoglobulin of interest wherein the cell contains a dominant negative allele of a mismatch repair gene. The properties of the immunoglobulin produced from the cells can be assayed to ascertain whether the immunoglobulin gene harbors a mutation. The assay may be directed to analyzing a polynucleotide encoding the immunoglobulin, or may be directed to the immunoglobulin polypeptide itself.

The invention also provides methods for generating a mutation in a gene affecting antibody production in an antibody-producing cell by culturing the cell expressing a dominant negative allele of a mismatch repair gene, and testing the cell to determine whether the cell harbors mutations within the gene of interest, such that a new biochemical feature (*e.g.*, over-expression and/or secretion of immunoglobulin products) is generated. The testing may include analysis of the steady state expression of the immunoglobulin gene of interest, and/or analysis of the amount of secreted protein encoded by the immunoglobulin gene of interest. The invention also embraces prokaryotic and eukaryotic transgenic cells made by this process, including cells from rodents, non-human primates and humans.

Other aspects of the invention encompass methods of reversibly altering the hypermutability of an antibody producing cell, in which an inducible vector containing a dominant negative allele of a mismatch repair gene operably linked to an inducible promoter is introduced into an antibody-producing cell. The cell is treated with an inducing agent to express the dominant negative mismatch repair gene (which can be *PMS2* (preferably human *PMS2*), *MLH1*, or *PMS1*). Alternatively, the cell may be induced to express a human *mutL* homolog or the first 133 amino acids of hPMS2. In another embodiment, the cells may be rendered capable of producing antibodies by co-transfecting a preselected immunoglobulin gene of interest. The immunoglobulin genes of the hypermutable cells, or the proteins produced by these methods may be analyzed for desired

properties, and induction may be stopped such that the genetic stability of the host cell is restored.

The invention also embraces methods of producing genetically altered antibodies by transfecting a polynucleotide encoding an immunoglobulin protein into a cell containing a dominant negative mismatch repair gene (either naturally or in which the dominant negative mismatch repair gene was introduced into the cell), culturing the cell to allow the immunoglobulin gene to become mutated and produce a mutant immunoglobulin, screening for a desirable property of said mutant immunoglobulin protein, isolating the polynucleotide molecule encoding the selected mutant immunoglobulin possessing the desired property, and transfecting said mutant polynucleotide into a genetically stable cell, such that the mutant antibody is consistently produced without further genetic alteration. The dominant negative mismatch repair gene may be *PMS2* (preferably human *PMS2*), *MLH1*, or *PMS1*. Alternatively, the cell may express a human *mutL* homolog or the first 133 amino acids of hPMS2.

The invention further provides methods for generating genetically altered cell lines that express enhanced amounts of an antigen binding polypeptide. These antigen-binding polypeptides may be, for example, immunoglobulins. The methods of the invention also include methods for generating genetically altered cell lines that secrete enhanced amounts of an antigen binding polypeptide. The cell lines are rendered hypermutable by dominant negative mismatch repair genes that provide an enhanced rate of genetic hypermutation in a cell producing antigen-binding polypeptides such as antibodies. Such cells include, but are not limited to hybridomas. Expression of enhanced amounts of antigen binding polypeptides may be through enhanced transcription or translation of the polynucleotides encoding the antigen binding polypeptides, or through the enhanced secretion of the antigen binding polypeptides, for example.

Methods are also provided for creating genetically altered antibodies *in vivo* by blocking the MMR activity of the cell host, or by transfecting genes encoding for immunoglobulin in a MMR defective cell host.

Antibodies with increased binding properties to an antigen due to genetic changes within the variable domain are provided in methods of the invention that block endogenous MMR of the cell host. Antibodies with increased binding properties to an antigen due to genetic changes within the CDR regions within the light and/or heavy

chains are also provided in methods of the invention that block endogenous MMR of the cell host.

The invention provides methods of creating genetically altered antibodies in MMR defective Ab producer cell lines with enhanced pharmacokinetic properties in host  
5 organisms including but not limited to rodents, primates, and man.

These and other aspects of the invention are provided by one or more of the embodiments described below. In one embodiment of the invention, a method for making an antibody producing cell line hypermutable is provided. A polynucleotide encoding a dominant negative allele of a MMR gene is introduced into an antibody-producing cell.  
10 The cell becomes hypermutable as a result of the introduction of the gene.

In another embodiment of the invention, a method is provided for introducing a mutation into an endogenous gene encoding for an immunoglobulin polypeptide or a single chain antibody. A polynucleotide encoding a dominant negative allele of a MMR gene is introduced into a cell. The cell becomes hypermutable as a result of the  
15 introduction and expression of the MMR gene allele. The cell further comprises an immunoglobulin gene of interest. The cell is grown and tested to determine whether the gene encoding for an immunoglobulin or a single chain antibody of interest harbors a mutation. In another aspect of the invention, the gene encoding the mutated immunoglobulin polypeptide or single chain antibody may be isolated and expressed in a  
20 genetically stable cell. In a preferred embodiment, the mutated antibody is screened for at least one desirable property such as, but not limited to, enhanced binding characteristics.

In another embodiment of the invention, a gene or set of genes encoding for Ig light and heavy chains or a combination therein are introduced into a mammalian cell host that is MMR defective. The cell is grown, and clones are analyzed for antibodies with  
25 enhanced binding characteristics.

In another embodiment of the invention, a method will be provided for producing new phenotypes of a cell. A polynucleotide encoding a dominant negative allele of a MMR gene is introduced into a cell. The cell becomes hypermutable as a result of the introduction of the gene. The cell is grown. The cell is tested for the expression of new  
30 phenotypes where the phenotype is enhanced secretion of a polypeptide.

These and other embodiments of the invention provide the art with methods that can generate enhanced mutability in cells and animals as well as providing cells and



animals harboring potentially useful mutations for the large-scale production of high affinity antibodies with beneficial pharmacokinetic profiles.

## BRIEF DESCRIPTION OF THE DRAWINGS

5           **Figure 1.** Hybridoma cells stably expressing PMS2 and PMS134 MMR genes. Shown is steady state mRNA expression of MMR genes transfected into a murine hybridoma cell line. Stable expression was found after 3 months of continuous growth. The (-) lanes represent negative controls where no reverse transcriptase was added, and the (+) lanes represent samples reverse transcribed and PCR amplified for the MMR  
10 genes and an internal housekeeping gene as a control.

**Figure 2.** Creation of genetically hypermutable hybridoma cells. Dominant negative MMR gene alleles were expressed in cells expressing a MMR-sensitive reporter gene. Dominant negative alleles such as PMS134 and the expression of MMR genes from other species results in antibody producer cells with a hypermutable phenotype that can be  
15 used to produce genetically altered immunoglobulin genes with enhanced biochemical features as well as lines with increased Ig expression and/or secretion. Values shown represent the amount of converted CPRG substrate which is reflective of the amount of function  $\beta$ -galactosidase contained within the cell from genetic alterations within the pCAR-OF reporter gene. Higher amounts of  $\beta$ -galactosidase activity reflect a higher  
20 mutation rate due to defective MMR.

**Figure 3.** Screening method for identifying antibody-producing cells containing antibodies with increased binding activity and/or increased expression/secretion

**Figure 4.** Generation of a genetically altered antibody with an increased binding activity. Shown are ELISA values from 96-well plates, screened for antibodies specific to  
25 hIgE. Two clones with a high binding value were found in HB134 cultures.

**Figure 5.** Sequence alteration within variable chain of an antibody (a mutation within the light chain variable region in MMR-defective HB134 antibody producer cells). Arrows indicate the nucleotide at which a mutation occurred in a subset of cells from a clone derived from HB134 cells. Panel A: The change results in a Thr to Ser  
30 change within the light chain variable region. The coding sequence is in the antisense direction. Panel B: The change results in a Pro to His change within the light chain variable region.

**Figure 6.** Generation of MMR-defective clones with enhanced steady state Ig

protein levels. A Western blot of heavy chain immunoglobulins from HB134 clones with high levels of MAb (>500ngs/ml) within the conditioned medium shows that a subset of clones express higher steady state levels of immunoglobulins (Ig). The H36 cell line was used as a control to measure steady state levels in the parental strain. Lane 1: fibroblast  
5 cells (negative control); Lane 2: H36 cell; Lane 3: HB134 clone with elevated MAb levels; Lane 4: HB134 clone with elevated MAb levels; Lane 5: HB134 clone with elevated MAb levels.

Methods have been discovered for developing hypermutable antibody-producing cells by taking advantage of the conserved mismatch repair (MMR) process of host cells.  
10 Dominant negative alleles of such genes, when introduced into cells or transgenic animals, increase the rate of spontaneous mutations by reducing the effectiveness of DNA repair and thereby render the cells or animals hypermutable. Hypermutable cells or animals can then be utilized to develop new mutations in a gene of interest. Blocking MMR in antibody-producing cells such as but not limited to: hybridomas; mammalian cells transfected with  
15 genes encoding for Ig light and heavy chains; mammalian cells transfected with genes encoding for single chain antibodies; eukaryotic cells transfected with Ig genes, can enhance the rate of mutation within these cells leading to clones that have enhanced antibody production and/or cells containing genetically altered antibodies with enhanced biochemical properties such as increased antigen binding. The process of MMR, also  
20 called mismatch proofreading, is carried out by protein complexes in cells ranging from bacteria to mammalian cells. A MMR gene is a gene that encodes for one of the proteins of such a mismatch repair complex. Although not wanting to be bound by any particular theory of mechanism of action, a MMR complex is believed to detect distortions of the DNA helix resulting from non-complementary pairing of nucleotide bases. The non-  
25 complementary base on the newer DNA strand is excised, and the excised base is replaced with the appropriate base, which is complementary to the older DNA strand. In this way, cells eliminate many mutations that occur as a result of mistakes in DNA replication.

Dominant negative alleles cause a MMR defective phenotype even in the presence of a wild-type allele in the same cell. An example of a dominant negative allele of a MMR  
30 gene is the human gene *hPMS2-134*, which carries a truncating mutation at codon 134 (SEQ ID NO:15). The mutation causes the product of this gene to abnormally terminate at the position of the 134th amino acid, resulting in a shortened polypeptide containing the N-terminal 133 amino acids. Such a mutation causes an increase in the rate of mutations,

which accumulate in cells after DNA replication. Expression of a dominant negative allele of a mismatch repair gene results in impairment of mismatch repair activity, even in the presence of the wild-type allele. Any allele which produces such effect can be used in this invention. Dominant negative alleles of a MMR gene can be obtained from the cells of

5 humans, animals, yeast, bacteria, or other organisms. Such alleles can be identified by screening cells for defective MMR activity. Cells from animals or humans with cancer can be screened for defective mismatch repair. Cells from colon cancer patients may be particularly useful. Genomic DNA, cDNA, or mRNA from any cell encoding a MMR protein can be analyzed for variations from the wild type sequence. Dominant negative

10 alleles of a MMR gene can also be created artificially, for example, by producing variants of the *hPMS2-134* allele or other MMR genes. Various techniques of site-directed mutagenesis can be used. The suitability of such alleles, whether natural or artificial, for use in generating hypermutable cells or animals can be evaluated by testing the mismatch repair activity caused by the allele in the presence of one or more wild-type alleles, to

15 determine if it is a dominant negative allele.

A cell or an animal into which a dominant negative allele of a mismatch repair gene has been introduced will become hypermutable. This means that the spontaneous mutation rate of such cells or animals is elevated compared to cells or animals without such alleles. The degree of elevation of the spontaneous mutation rate can be at least 2-

20 fold, 5-fold, 10-fold, 20-fold, 50-fold, 100-fold, 200-fold, 500-fold, or 1000-fold that of the normal cell or animal. The use of chemical mutagens such as but limited to methane sulfonate, dimethyl sulfonate, O6-methyl benzadine, MNU, ENU, etc. can be used in MMR defective cells to increase the rates an additional 10 to 100 fold that of the MMR deficiency itself.

25 According to one aspect of the invention, a polynucleotide encoding for a dominant negative form of a MMR protein is introduced into a cell. The gene can be any dominant negative allele encoding a protein, which is part of a MMR complex, for example, *PMS2*, *PMS1*, *MLH1*, or *MSH2*. The dominant negative allele can be naturally occurring or made in the laboratory. The polynucleotide can be in the form of genomic DNA, cDNA, RNA,

30 or a chemically synthesized polynucleotide.

The polynucleotide can be cloned into an expression vector containing a constitutively active promoter segment (such as but not limited to CMV, SV40, Elongation Factor or LTR sequences) or to inducible promoter sequences such as the steroid inducible

pIND vector (Invitrogen), where the expression of the dominant negative MMR gene can be regulated. The polynucleotide can be introduced into the cell by transfection.

According to another aspect of the invention, an immunoglobulin (Ig) gene, a set of Ig genes or a chimeric gene containing whole or parts of an Ig gene can be transfected into MMR deficient cell hosts, the cell is grown and screened for clones containing genetically altered Ig genes with new biochemical features. MMR defective cells may be of human, primates, mammals, rodent, plant, yeast or of the prokaryotic kingdom. The mutated gene encoding the Ig with new biochemical features may be isolated from the respective clones and introduced into genetically stable cells (*i.e.*, cells with normal MMR) to provide clones that consistently produce Ig with the new biochemical features. The method of isolating the Ig gene encoding Ig with new biochemical features may be any method known in the art. Introduction of the isolated polynucleotide encoding the Ig with new biochemical features may also be performed using any method known in the art, including, but not limited to transfection of an expression vector containing the polynucleotide encoding the Ig with new biochemical features. As an alternative to transfecting an Ig gene, a set of Ig genes or a chimeric gene containing whole or parts of an Ig gene into an MMR deficient host cell, such Ig genes may be transfected simultaneously with a gene encoding a dominant negative mismatch repair gene into a genetically stable cell to render the cell hypermutable.

Transfection is any process whereby a polynucleotide is introduced into a cell. The process of transfection can be carried out in a living animal, *e.g.*, using a vector for gene therapy, or it can be carried out *in vitro*, *e.g.*, using a suspension of one or more isolated cells in culture. The cell can be any type of eukaryotic cell, including, for example, cells isolated from humans or other primates, mammals or other vertebrates, invertebrates, and single celled organisms such as protozoa, yeast, or bacteria.

In general, transfection will be carried out using a suspension of cells, or a single cell, but other methods can also be applied as long as a sufficient fraction of the treated cells or tissue incorporates the polynucleotide so as to allow transfected cells to be grown and utilized. The protein product of the polynucleotide may be transiently or stably expressed in the cell. Techniques for transfection are well known. Available techniques for introducing polynucleotides include but are not limited to electroporation, transduction, cell fusion, the use of calcium chloride, and packaging of the polynucleotide together with lipid for fusion with the cells of interest. Once a cell has been transfected with the MMR

gene, the cell can be grown and reproduced in culture. If the transfection is stable, such that the gene is expressed at a consistent level for many cell generations, then a cell line results.

An isolated cell is a cell obtained from a tissue of humans or animals by  
5 mechanically separating out individual cells and transferring them to a suitable cell culture medium, either with or without pretreatment of the tissue with enzymes, *e.g.*, collagenase or trypsin. Such isolated cells are typically cultured in the absence of other types of cells. Cells selected for the introduction of a dominant negative allele of a mismatch repair gene may be derived from a eukaryotic organism in the form of a primary cell culture or an  
10 immortalized cell line, or may be derived from suspensions of single-celled organisms.

A polynucleotide encoding for a dominant negative form of a MMR protein can be introduced into the genome of an animal by producing a transgenic animal. The animal can be any species for which suitable techniques are available to produce transgenic animals. For example, transgenic animals can be prepared from domestic livestock, *e.g.*,  
15 bovine, swine, sheep, goats, horses, etc.; from animals used for the production of recombinant proteins, *e.g.*, bovine, swine, or goats that express a recombinant polypeptide in their milk; or experimental animals for research or product testing, *e.g.*, mice, rats, guinea pigs, hamsters, rabbits, etc. Cell lines that are determined to be MMR defective can then be used as a source for producing genetically altered immunoglobulin genes *in vitro*  
20 by introducing whole, intact immunoglobulin genes and/or chimeric genes encoding for single chain antibodies into MMR defective cells from any tissue of the MMR defective animal.

Once a transfected cell line or a colony of transgenic animals has been produced, it can be used to generate new mutations in one or more gene(s) of interest. A gene of  
25 interest can be any gene naturally possessed by the cell line or transgenic animal or introduced into the cell line or transgenic animal. An advantage of using such cells or animals to induce mutations is that the cell or animal need not be exposed to mutagenic chemicals or radiation, which may have secondary harmful effects, both on the object of the exposure and on the workers. However, chemical mutagens may be used in  
30 combination with MMR deficiency, which renders such mutagens less toxic due to an undetermined mechanism. Hypermutable animals can then be bred and selected for those producing genetically variable B-cells that may be isolated and cloned to identify new cell lines that are useful for producing genetically variable cells. Once a new trait is identified,

the dominant negative MMR gene allele can be removed by directly knocking out the allele by technologies used by those skilled in the art or by breeding to mates lacking the dominant negative allele to select for offspring with a desired trait and a stable genome.

Another alternative is to use a CRE-LOX expression system, whereby the dominant

- 5 negative allele is spliced from the animal genome once an animal containing a genetically diverse immunoglobulin profile has been established. Yet another alternative is the use of inducible vectors such as the steroid induced pIND (Invitrogen) or pMAM (Clonotech) vectors which express exogenous genes in the presence of corticosteroids.

- 10 Mutations can be detected by analyzing for alterations in the genotype of the cells or animals, for example by examining the sequence of genomic DNA, cDNA, messenger RNA, or amino acids associated with the gene of interest. Mutations can also be detected by screening for the production of antibody titers. A mutant polypeptide can be detected by identifying alterations in electrophoretic mobility, spectroscopic properties, or other physical or structural characteristics of a protein encoded by a mutant gene. One can also
- 15 screen for altered function of the protein *in situ*, in isolated form, or in model systems. One can screen for alteration of any property of the cell or animal associated with the function of the gene of interest, such as but not limited to Ig secretion.

Examples of mismatch repair proteins and nucleic acid sequences include the following:

20

#### PMS2 (mouse) (SEQ ID NO:5)

- |    |            |             |             |            |            |            |     |
|----|------------|-------------|-------------|------------|------------|------------|-----|
|    | MEQTEGVSTE | CAKAIKPIDG  | KSVHQICSGQ  | VILSLSTAVK | ELIENSVDAG | ATTIDLRLKD | 60  |
|    | YGVDLIEVSD | NGCGVEEENF  | EGLALKHHTS  | KIQEFADLTQ | VETFGFRGEA | LSSLCALSDV | 120 |
|    | TISTCHGSAS | VGTRLVFDHN  | GKITQKTPYP  | RPKGTTVSVQ | HLFYTLFVRY | KEFQRNIKKE | 180 |
| 25 | YSKMVQVLQA | YCIIISAGVRV | SCTNQLGQ GK | RHAVVCTSGT | SGMKENIGSV | FGQKQLQSLI | 240 |
|    | PFVQLPPSDA | VCEEYGLSTS  | GRHKTFTSTF  | ASFHSARTAP | GGVQQTGSFS | SSIRGPVTQQ | 300 |
|    | RSLSLSMRFY | HMYNRHQYPF  | VVLNVSVDS   | CVDINVTDPK | RQILLQEEKL | LLAVLKTSLI | 360 |
|    | GMFDSANKL  | NVNQQPLLDV  | EGNLVKLHTA  | ELEKPVPGKQ | DNSPSLKSTA | DEKRVASISR | 420 |
|    | LREAFSLHPT | KEIKSRGPET  | AELTRSFPS   | KRGVLSSYPS | DVISYRGLRG | SQDKLVSPTD | 480 |
| 30 | SPGDCMDREK | IEKDSGLSST  | SAGSEEEFST  | PEVASSFSSD | YNVSSLEDRP | SQETINCSDL | 540 |
|    | DCRPPGTGQS | LKPEDHGYQC  | KALPLARLSP  | TNAKRFKTEE | RPSNVNISQR | LPGPOSTSAA | 600 |
|    | EVDVAIKMNK | RIVLLEFSL   | SLAKRMKQLQ  | HLKAQNKHEL | SYRKFRKIC  | PGENQAAEDE | 660 |
|    | LRKEISKSMF | AEMEILGQFN  | LGFIIVTKLE  | DLFLVDQHAA | DEKYNFEMLQ | QHTVLAQRL  | 720 |
|    | ITPQTLNLTA | VNEAVLIENL  | EIFRKNGFDF  | VIDEDAPVTE | RAKLISLPTS | KNWTFGPQDI | 780 |
| 35 | DELIFMLSDS | PGVMCRPSRV  | RQMFASRACR  | KSVMIGTALN | ASEMKKLITH | MGEMDHPWNC | 840 |
|    | PHGRPTMRHV | ANLDVISQN   |             |            |            |            | 859 |

#### PMS2 (mouse cDNA) (SEQ ID NO:6)

- |    |            |            |            |             |            |            |     |
|----|------------|------------|------------|-------------|------------|------------|-----|
| 40 | gaattccggt | gaaggtcctg | aagaatttcc | agattcctga  | gtatcattgg | aggagacaga | 60  |
|    | taacctgtcg | tcaggtaacg | atggtgtata | tgcaacagaa  | atgggtgttc | ctggagacgc | 120 |
|    | gtcttttccc | gagagcggca | ccgcaactct | cccgcggtga  | ctgtgactgg | aggagtcctg | 180 |
|    | catccatgga | gcaaaccgaa | ggcgtgagta | cagaatgtgc  | taaggccatc | aagcctattg | 240 |
|    | atgggaagtc | agtccatcaa | atttggtctg | ggcagggtgat | actcagttta | agcaccgctg | 300 |
| 45 | tgaaggagtt | gatagaaaat | agtgtagatg | ctgggtgctac | tactattgat | ctaaggctta | 360 |

	aagactatgg	ggtggacctc	attgaagttt	cagacaatgg	atgtggggta	gaagaagaaa	420
	actttgaagg	tctagctctg	aaacatcaca	catctaagat	tcaagagttt	gccgacctca	480
	cgcagggttg	aactttcggc	tttcgggggg	aagctctgag	ctctctgtgt	gcactaagtg	540
5	atgtcactat	atctacctgc	cacgggtctg	caagcggttg	gactcgactg	gtgtttgacc	600
	ataatgggaa	aatcaccocag	aaaactccct	accccggacc	taaagggaacc	acagtcagtg	660
	tgcagcactt	atcttataca	ctacccgtgc	gttacaaga	gtttcagagg	aacattaaaa	720
	aggagtattc	caaaatgggtg	caggtcttac	aggcgtagtg	tatcatctca	gcaggcgctcc	780
	gtgtaagctg	cactaatcag	ctcggacagg	ggaagcgga	cgctgtggtg	tgacaagcg	840
	gcacgtctgg	catgaaggaa	aatatcgggt	ctgtgtttgg	ccagaagcag	ttgcaaagcc	900
10	tcattccttt	tggtcagctg	ccccctagtg	acgctgtgtg	tgaagagtac	ggcctgagca	960
	cttcaggagc	ccacaaaacc	ttttctacgt	ttcgggcttc	atctcacagt	gcacgcaagg	1020
	cgcggggagg	agtgaacag	acaggcagtt	tttcttcac	aatcagaggc	cctgtgaccc	1080
	agcaaaggtc	tctaagcttg	tcaatgaggt	tttatcacat	gtataaccgg	catcagtacc	1140
	cattttgtcgt	ccttaacgtt	tcogttgact	cagaatgtgt	ggatattaat	gtaactccag	1200
15	ataaaaggca	aattctacta	caagaagaga	agctattgct	ggcgttttta	aagacctcct	1260
	tgataggaat	gtttgacagt	gatgcaaca	agcttaagt	caaccagcag	ccactgctag	1320
	atgttgagg	taacttagta	aagctgcata	ctgcagaact	agaaaagcct	gtgccaggaa	1380
	agcaagataa	ctctccttca	ctgaagagca	cagcagacga	gaaaagggtg	gcatccatct	1440
20	ccaggctgag	agaggccttt	tctcttcac	ctactaaaga	gatcaagtct	aggggtccag	1500
	agactgctga	actgacacgg	agttttccaa	gtgagaaaag	ggcgctgtta	tcctcttctc	1560
	cttcagacgt	catctcttac	agaggcctcc	gtggctcgca	ggacaaattg	gtgagtccca	1620
	cggacagccc	tgggtagctg	atggacagag	tgaaaataga	aaaagactca	gggctcagca	1680
	gcacctcagc	tggctctgag	gaagagttca	gcaccccgca	agtggccagt	agcttttagca	1740
25	gtgactataa	cgtgagctcc	ctagaagaca	gaccttctca	ggaaaccata	aactgtggtg	1800
	acctggactg	ccgtcctcca	ggtacaggac	agtcttgaa	gccagaagac	catggatata	1860
	aatgcaaagc	tctacctcta	gctcgtctgt	caccacaaa	tgccaagcgc	ttcaagacag	1920
	aggaaagacc	ctcaaagtgc	aacattttct	aaagattgcc	tggtcctcag	agcacctcag	1980
	cagctgaggt	cgatgtagcc	ataaaaatga	ataagagaat	cgtgctcctc	gagttctctc	2040
30	tgagttctct	agctaagcga	atgaagcagt	tacagcacct	aaaggcgag	aacaaacatg	2100
	aactgagtta	cagaaaat	agggccaaga	tttgccctgg	agaaaaccaa	gcagcagaag	2160
	atgaactcag	aaaagagatt	agtaaatcga	tggttgagca	gatggagatc	ttgggtcagt	2220
	ttaacctggg	atcttatagta	accaaactga	aagaggacct	cttctggtg	gaccagcatg	2280
35	ctgocgatga	gaagtacaac	tttgagatgc	tgacgagca	cacggtgctc	caggcgagaa	2340
	ggctcatcac	accccagact	ctgaacttaa	ctgctgtcaa	tgaagctgta	ctgatgagga	2400
	atctggaaat	attcagaaaag	aatggccttg	actttgtcat	tgatgaggat	gctccagtca	2460
	ctgaaagggc	taaattgatt	tccttaccaa	ctagtaaaaa	ctggaccttt	ggaccccaag	2520
	atatagatga	actgatcttt	atgttaagt	acagccctgg	ggatcatgtg	cggccctcac	2580
	gagtcagaca	gatgtttgct	tcagagcct	gtcgaagtc	agtgatgatt	ggaacggcgc	2640
40	tcaatgcgag	cgagatgaag	aagctcatca	cccacatggg	tgagatggac	cacccttgga	2700
	actgccccca	cggcaggcca	accatgaggc	acgttgccaa	tctggatgtc	atctctcaga	2760
	actgacacac	ccctttagtc	atagagttta	ttacagattg	ttcggtttgc	aaagagaagg	2820
	tttttaagtaa	tctgattatc	gttgtagaaa	aattagcatg	ctgctttaat	gtactggatc	2880
	cattttaaag	cagtgttaag	gcaggcatga	tggagtggtc	ctctagctca	gctacttggg	2940
45	tgatccgggt	ggagctcatg	tgagcccagg	actttgagac	cactccgagc	cacattcatg	3000
	agactcaatt	caaggacaaa	aaaaaaaaa	tatttttgaa	gccttttaaa	aaaaaa	3056

## PMS2 (human) (SEQ ID NO:7)

	MERAESSSTE	PAKAIKPIDR	KSVHQICSGQ	VVLSLSTAVK	ELVENSIDAG	ATNIDLKLKD	60
50	YGVDLIEVSD	NGCGVEEENF	EGLTLKHHTS	KIQEFADLTQ	VETFGFRGEA	LSSLCALSDV	120
	TISTCHASAK	VGTRLMFDHN	GKIIQKTPYP	RPRGTTVSQ	QLFSTLPVRH	KEFQRNIKKE	180
	YAKMVQVLHA	YCIISAGIRV	SCTNQLGQK	RQPVVCTGGS	PSIKENIGSV	FGQKQLQSLI	240
	PFVQLPPSDS	VCEEYGLSCS	DALHNLFYIS	GFISQCTHGV	GRSSTDROFF	FINRRPCDPA	300
	KVCRLVNEVY	HMYNRHQYPF	VVLNISVDSE	CVDINVTPDK	RQILLQEEKL	LLAVLKTSLI	360
55	GMFSDSVNKL	NVSQQPLLDV	EGNLIKMHAA	DLEKPMVEKQ	DQSPSLRTGE	EKKDVSISRL	420
	REAFSLRHTT	ENKPHSPKTP	EPRRSPLGQK	RGMSSSTSG	AISDKGVLRP	QKEAVSSSHG	480
	PSDPTDRAEV	EKDSGHGSTS	VDSEGFISIP	TGSHCSSEYA	ASSPGDRGSQ	EHVDSQEKAP	540
	ETDDSFSDVD	CHSNQEDTGC	KFRVLPQPTN	LATPNTKRFK	KEEILSSSDI	CQKLVTQDM	600
	SASQVDVAVK	INKKVPLDF	SMSSLAKRIK	QLHHEAQOSE	GEQNYRKFR	KICPGENQAA	660
60	EDELIRKEISK	TMFAEMEITG	QFNLFITITK	LNEDIFIVDQ	HATDEKYNFE	MLQOHTVLQG	720
	QRLIAPQTLN	LTAVNEAVLI	ENLEIFRKN	FDFVIDENAP	VTERAKLISL	PTSKNWTFGP	780
	QDVDELIFML	SDSPGVMCRP	SRVKQMFASR	ACRKSVMIGT	ALNTSEMKKL	ITHMGEMDHP	840
	WNCPHGRPTM	RHIANLGVIS	QN				862

**PMS2 (human cDNA) (SEQ ID NO:8)**

	cgaggcgggat	cggtgtgtgc	atccatggag	cgagctgaga	gctcgagtag	agaacctgct	60
	aaggccatca	aacctattga	tcggaagtca	gtccatcaga	tttgctctgg	gcaggtggta	120
	ctgagctctaa	gcactgcggt	aaaggagtta	gtagaaaaca	gtctggatgc	tggtgccact	180
5	aatattgatc	taaagcttaa	ggactatgga	gtggatctta	ttgaagtttc	agacaatgga	240
	tgtggggtag	aagaagaaaa	cttcgaaggc	ttaactctga	aacatcacac	atctaagatt	300
	caagagtttg	ccgacctaac	tcaggttgaa	acttttggct	ttcgggggga	agctctgagc	360
	tcactttgtg	cactgagcga	tgtcaccatt	tctacctgcc	acgcacgcgc	gaagggttga	420
	actcgactga	tgtttgatca	caatgggaaa	attatccaga	aaaccccccta	ccccgcgcc	480
10	agagggacca	cagtcagcgt	gcagcagtta	ttttccacac	tacctgtgcg	ccataaggaa	540
	tttcaaagga	atattaagaa	ggagtatgcc	aaaatgggtcc	aggtcttaca	tgcatactgt	600
	atcattttcag	caggcatccg	tgttaagttgc	accaatcagc	ttggacaagg	aaaacgacag	660
	cctgtgggtat	gcacaggtgg	aagccccagc	ataaaggaaa	atatcggtgc	tgtgtttggg	720
	cagaagcaggt	tgcaaacgct	cattccctttt	gttcagctgc	cccctagtga	ctccgtgtgt	780
15	gaagagtagc	gtttgagctg	ttcggatgct	ctgcataatc	ttttttacat	ctcaggtttc	840
	atttcacaat	gcacgcatgg	agttggaagg	agttcaacag	acagacagtt	tttctttatc	900
	aaccggcggc	cttgtgaccc	agcaaaggtc	tgcagactcg	tgaatgaggt	ctaccacatg	960
	tataatcgac	accagtatcc	atttgttgtt	cttaacattt	ctgttgatcc	agaatgcgtt	1020
	gataccaatg	ttactccaga	taaaaggcaa	attttgctac	aagaggaaaa	gcttttggttg	1080
20	gcagtttttaa	agacctcttt	gataggaatg	tttgatagtg	atgtcaacaa	gctaaatgtc	1140
	agtcagcagc	cactgctgga	tgttgaaagt	aacttaataa	aatgcatgc	agcggatttg	1200
	gaaaagccca	tggtagaaaa	gcaggatcaa	tccccttcat	taaggactgg	agaagaaaaa	1260
	aaagacgtgt	ccatttccag	actgcgagag	gcctttttctc	ttcgtcacac	aacagagaac	1320
	aagcctcaca	gcccacagac	tccagaaaca	agaaggagcc	ctctaggaca	gaaaaggggt	1380
25	atgctgtctt	ctagacttcc	aggtgccatc	tctgacaaag	gcgtcctgag	acctcagaaa	1440
	gaggcagtag	gttccagtag	cggacccagt	gaccctacgg	acagagcggg	gggtggagaag	1500
	gactcggggc	acggcagcac	ttccgtggat	tctgaggggt	tcagcatccc	agacacgggc	1560
	agtcactgca	gcagcgagta	tgcggccagc	tccccagggg	acaggggctc	gcaggaacat	1620
	gtggactctc	aggagaaaag	gcctgaaact	gaogactctt	tttcagatgt	ggactgccat	1680
30	tcaaacccag	aagataccgg	atgtaaattt	cgagtttttg	ctcagccaac	taattctgca	1740
	accccaaaaa	caaacggttt	taaaaaagaa	gaaattcttt	ccagttctga	caatttgtcaa	1800
	aagttagtaa	atactcagga	catgtcagcc	tctcaggttg	atgtagctgt	gaaaattaat	1860
	aagaaagttag	tgccccctgga	cttttctatg	agttcttttag	ctaaacgaat	aaagcagtta	1920
	catcatgaag	cacagcaaaag	tgaaggggaa	cagaattaca	ggaagttagg	ggcaagattt	1980
35	tgtcctggag	aaaaatcaagc	agccgaagat	gaactaagaa	aagagataag	taaaacgatg	2040
	tttgacagaaa	tggaaatcat	tggtcagttt	aacctgggat	ttataataac	caaacatgaat	2100
	gaggatatct	tcatagtgga	ccagcatgcc	acggacgaga	agtataactt	cgagatgctg	2160
	cagcagcaca	ccgtgctcca	ggggcagagg	ctcatagcac	ctcagactct	caacttaact	2220
	gctgttaaatg	aagctgttct	gatagaaaat	ctggaaatat	ttagaaagaa	tggtctttgat	2280
40	tttgtttatcg	atgaaaatgc	tccagtcact	gaaagggcta	aactgatttc	cttgccaact	2340
	agtaaaaaat	ggaccttcgg	acccaggagc	gtcagtgaa	tgatcttcat	gctgagcgac	2400
	agccctgggg	tcagtgtccg	gccttcccg	gtcaagcaga	tggttgccctc	cagagcctgc	2460
	cggaagtccg	tgatgattgg	gactgctctt	aacacaagcg	agatgaagaa	actgatcacc	2520
	cacatggggg	agatggacca	cccctggaac	tgtcccatg	gaaggccaac	catgagacac	2580
45	atcgccaacc	tgggtgtcat	ttctcagaac	tgaccgtagt	cactgtatgg	aataattggg	2640
	tttatcgtag	atttttatgt	tttgaaagac	agagtcttca	ctaacctttt	ttgttttaaa	2700
	atgaaacctg	ctacttaaaa	aaaatacaca	tcacacccat	ttaaaagtga	tcttgagaac	2760
	cttttcaaac	c					2771

**PMS1 (human) (SEQ ID NO:9)**

50	MKQLPAATVR	LLSSSQIITS	VSVVKELIE	NSLDAGATSV	DVKLENYGFD	KIEVRDNGEG	60
	IKAVDAPVMA	MKYYSKINS	HEDLENLTYY	GFRGEALGSI	CCIAEVLITT	RTAADNFSTQ	120
	YVLDGSGHIL	SQKPSHLGQG	TTVTALRLFK	NLPVRKQFYS	TAKKCKDEIK	KIQDLLMSFG	180
	ILKPDRLRIVF	VHNKAVIWQK	SRVSDHKMAL	MSVLGTAVMN	NMESFQYHSE	ESQIYLSGFL	240
55	PKCDADHSFT	SLSTPERSFI	FINSRPVHQK	DILKLIRHHY	NLKCLKESTR	LYPVFFLKID	300
	VPTADVVDNL	TPDKSQVLLQ	NKESVLIALE	NLMTTCYGPL	PSTNSYENNK	TDVSAADIVL	360
	SKTAETDVLF	NKVESSGKNY	SNVDTSVIPF	QNDMHNDESG	KNTDDCLNHQ	ISIGDFGYGH	420
	CSSEISNIDK	NTKNAFQDIS	MSNVSWENSQ	TEYSKTCFIS	SVKHTQSENG	NKDHIDESGE	480
	NEEEAGLENS	SEISADEWSR	GNILKNSVGE	NIEPVKILVP	EKSLPCKVSN	NNYPIPEQMN	540
60	LNEDSCNKKK	NVIDNKSCKV	TAYDLLSNRV	IKKEMSASAL	FVQDHRPQFL	IENPKTSLED	600
	ATLQIEELWK	TLSEEEKLKY	EEKATKDLER	YNSQMKRAIE	QESQMSLKDG	RKKIKPTSAS	660
	NLAQKHKLKT	SLSNQPKLDE	LLQSQIEKRR	SQNIKMVQIP	FSMKNLKINF	KKQNKVDLEE	720
	KDEPCLIHNL	RFPDAWLMTS	KTEVMLLNPY	RVEEALLFKR	LENHKLPAE	PLEKPIMLTE	780
	SLFNGSHYLD	VLYKMTADDQ	RYSGSTYLS	PRLTANGFKI	KLIPGVSITE	NYLEIEGMAN	840
65	CLPFYGVADL	KEILNAILNR	NAKEVYECRP	RKVISYLEGE	AVRLSRQLPM	YLSKEDIQDI	900
	IYRMKHQFGN	EIKECVHGRP	FFHHLTYLPE	TT			932



**PMS1 (human) (SEQ ID NO:10)**

	ggcaogagtg	gctgcttgcg	gctagtggtg	ggtaattgcc	tgcctcgccg	tagcagcaag	60
	ctgctctgtt	aaaagcgaaa	atgaaacaat	tgctcgccgc	aacagttcga	ctcctttcaa	120
5	gttctcagat	catcacttcg	gtggctcagtg	ttgtaaaaga	gcttattgaa	aaotccttgg	180
	atgctgggtgc	cacaagcgta	gatgttaaac	tggagaacta	tggatttgat	aaaattgagg	240
	tgcgagataa	cggggagggg	atcaaggctg	tggatgcacc	tgtaatggca	atgaagtact	300
	acacctcaaa	aataaatagt	catgaagatc	ttgaaaattt	gacaacttac	ggttttctgt	360
	gagaagcctt	ggggccaatt	tgttgatatag	ctgaggtttt	aattacaaca	agaacggctg	420
10	ctgataattt	tagcaccag	tatgttttag	atggcagtg	ccacatactt	tctcagaaac	480
	cttcacatct	tggtcaaggt	acaactgtaa	ctgctttaag	attattttaag	aatctacctg	540
	taagaaagca	gttttactca	actgcáaaaa	aatgtaaaga	tgaaataaaa	aagatccaag	600
	atctcctcat	gagctttggt	atccttaaac	ctgacttaag	gattgtcttt	gtacataaca	660
	aggcagttat	ttggcagaaa	agcagagtat	cagatcacia	gatggctctc	atgtcagttc	720
15	tggggaactgc	tgttatgaac	aatatggaat	cctttcagta	ccactctgaa	gaatctcaga	780
	tttatctcag	tggattttct	ccaaagtgtg	atgcagacca	ctctttcact	agtctttcaa	840
	caccagaaag	aagtttcatc	ttcataaaca	gtcgaccagt	acatcaaaaa	gatatcttaa	900
	agttaatccg	acatcattac	aatctgaaat	gcctaaagga	atctactcgt	ttgtatcctg	960
	ttttctttct	gaaaatcgat	gttctctacag	ctgatgttga	tgtaaattta	acaccagata	1020
20	aaagccaagt	attattacaa	aataaggaat	ctgttttaat	tgctcttgaa	aatctgatga	1080
	cgacttggtta	tggaccattta	cctagtagtaa	attcttatga	aaataataaaa	acagatgttt	1140
	ccgcagctga	catcgttctt	agtaaaacag	cagaaacaga	tgtgcttttt	aataaagtgg	1200
	aatcatctgg	aaagaattat	tcaaagtgtt	atacttcagt	cattccattc	caaaatgata	1260
	tgcataatga	tgaatctgga	aaaaacactg	atgattgttt	aaatcaccag	ataagtattg	1320
25	gtgacttttg	ttatggatcat	tgtagtagtg	aaatttctaa	cattgataaa	aacactaaga	1380
	atgcatttca	ggacatttca	atgagtaatg	tatcatggga	gaactctcag	acggaatata	1440
	gtaaaaacttg	ttttataagt	tccgttaagc	acaccagtc	agaaaaatggc	aataaagacc	1500
	atatagatga	gagtggggaa	aatgaggaag	aagcaggtct	tgaaaactct	tcggaaattt	1560
	ctgcagatga	gtggagcagg	ggaaatatac	ttaaaaattc	agtgggagag	aatattgaa	1620
30	ctgtgaaaat	tttagtgctt	gaaaaaagtt	taccatgtaa	agtaagtatt	aataattatc	1680
	caatccctga	acaaatgaat	cttaatgaag	attcatgtaa	caaaaaatca	aatgtaatag	1740
	ataataaatc	tggaaaagtt	acagcttatg	atttacttag	caatcgagta	atcaagaaac	1800
	ccatgtcagc	aagtgtctct	tttgttcaag	atcatcgctc	tcagtttctc	atagaaaatc	1860
	ctaagactag	tttagaggat	gcaacactac	aaattgaaga	actgtggaag	acattgagtg	1920
35	aagaggaaaa	actgaaatat	gaagagaagg	ctactaaaga	ccttggaacga	tacaatagtc	1980
	aaatgaagag	agccattgaa	caggagtcac	aaatgtcact	aaaagatggc	agaaaaaaga	2040
	taaaacccac	cagcgcatgg	aatttggtcc	agaagcacia	gttaaaaacc	tcattatcta	2100
	atcaaccaaa	acttgatgaa	ctccttcagt	cccaaattga	aaaaagaagg	agtcaaaaata	2160
	ttaaaatggt	acagatcccc	ttttctatga	aaaacttaaa	aataaatttt	aagaaacaaa	2220
40	acaaagttag	cttagaagag	aaggatgaac	cctgcttgat	ccacaatctc	agggttcctg	2280
	atgcatggct	aatgacatcc	aaaacagagg	taatgttatt	aaatccatat	agagtagaag	2340
	aagccctgct	attttaaaga	cttcttgaga	atcataaaact	tcctgcagag	ccactggaaa	2400
	agccaattat	gttaacagag	agtcttttta	atggatctca	ttatttagac	gttttatata	2460
	aatgacagc	agatgaccaa	agatacagtg	gatcaactta	cctgtctgat	cctcgtctta	2520
45	cagcgaatgg	tttcaagata	aaattgatac	caggagtttc	aattactgaa	aattacttgg	2580
	aaatagaagg	aatggctaatt	tgtctcccat	tctatggagt	agcagattta	agcaaaattc	2640
	ttaatgctat	attaaacaga	aatgcaaagg	aagtttatga	atgtagacct	cgcaaagtga	2700
	taagttattt	agaggagaa	gcagtgcgtc	tatccagaca	attacccatg	tacttatcaa	2760
	aagaggacat	ccaagacatt	atctacagaa	tgaagcacca	gtttggaaat	gaaattaaag	2820
50	agtgtgttca	tggtcgccca	ttttttcatc	atttaacctt	tcttccagaa	actacatgat	2880
	taaatatggt	taagaagatt	agttaccatt	gaaattgggt	ctgtcataaa	acagcatgag	2940
	tctgggttta	aattatcttt	gtattatgtg	tcacatgggt	atttttttaa	tgaggattca	3000
	ctgacttggt	tttatattga	aaaaagttcc	acgtattgta	gaaaacgtaa	ataaactaat	3060
55	aac						3063

**MSH2 (human) (SEQ ID NO:11)**

60	MAVQPKETLQ	LESAAEVGFV	RFFQGMPEKP	TTTVRLFDRG	DFYTAHGEDA	LLAAREVFKE	60
	QGVIKYMGPA	GAKNLQSVVL	SKMNFESFVK	DLLLVRQYRV	EVYKNRAGNK	ASKENDWYLA	120
	YKASPGNLSQ	FEDILFGNND	MSASIGVGVV	KMSAVDQGRQ	VGVGVDVSIQ	RKLGCEFPD	180
	NDQFSNLEAL	LIQIGPKECV	LPGGETAGDM	GKLRQIIQRG	GILITERKKA	DFSTKDIYQD	240
	LNRLKKGKKG	EQMNSAVLPE	MENQVAVSSL	SAVIKFLLELL	SDDSNFGQFE	LTTDFDSQYM	300
65	KLDIAAVRAL	NLFQGSVEDT	TGSQSLAALL	NKCKTPQGQR	LVNQWIKQPL	MDKNRIEERL	360

	NLVEAFVEDA	ELRQTLQEDL	LRFPDLNRL	AKKFQROAAN	LQDCYRLYQG	INQLPNVIOA	420
	LEKHEGKHQK	LLAVFVTPL	TDLRSDFSKF	QEMIETTLDM	DQVENHEFLV	KPSFDPNLSE	480
	LREIMNDLEK	KMQSTLISAA	RDLGLDPGKQ	IKLDSSAQFG	YYFRVTCKEE	KVLRNNKNFS	540
	TVDIQKNGVK	FTNSKLTSLN	EEYTKNKTEY	EEAQDAIVKE	IVNISSGYVE	PMQTLNDVLA	600
5	QLDAVVSFAH	VSNGAPVPYV	RPAILEKGQG	RIILKASRHA	CVEVQDEIAF	IPNDVYFEKD	660
	KQMFHIIITGP	NMGGKSTYIR	QTGVIVLMAQ	IGCFVPCESA	EVSIVDCILA	RVGAGDSQLK	720
	GVSTFMAEML	ETASILRSAT	KDSLIIIDEL	GRGTSTYDGF	GLAWAISEYI	ATKIGAFCMF	780
	ATHFHELTAL	ANQIPTVNNL	HVTALTTEET	LTMLYQVKKG	VCDQSFGIHV	AELANFPKHV	840
	IECAKQKALE	LEEFQYIGES	QGYDIMEPAA	KKCYLEREQG	EKIIQEFLSK	VKQMPFTEMS	900
10	EENITIKLKQ	LKAEVIAKNN	SFVNEIISRI	KVTT			934

## MSH2 (human cDNA) (SEQ ID NO:12)

	ggcggaac	agcttagtgg	gtgtggggtc	gcgcattttc	ttcaaccagg	aggtgaggag	60
	gtttcgacat	ggcgggtgcag	ccgaaggaga	cgctgcagtt	ggagagcgcg	gccgaggctcg	120
15	gcttcgtgcg	cttctttcag	ggcatgccgg	agaagccgac	caccacagtg	cgccttttcg	180
	accggggcga	cttctatacg	gcgcacggcg	aggacgcgct	gctggccgcc	cgggagctgt	240
	tcaagaccca	gggggtgatc	aagtacatgg	ggccggcagg	agcaaagaat	ctgcagagtg	300
	ttgtgcttag	taaaatgaat	tttgaatctt	ttgtaaaaga	tcttctctcg	gttcgctcagt	360
	atagagttga	agtttataag	aatagagctg	gaaataaggc	atccaaggag	aatgatttgt	420
20	atttggcata	taaggcttct	cctggcaatc	tctctcagtt	tgaagacatt	ctctttggta	480
	acaatgatat	gtcagcttcc	attgggtgtg	tgggtgttaa	aatgtccgca	gttgatggcc	540
	agagacaggt	tggagttggg	tatgtggatt	ccatacagag	gaaactagga	ctgtgtgaat	600
	tccctgataa	tgatcagttc	tccaatcttg	aggctctcct	catccagatt	ggaccaaagg	660
	aatgtgtttt	accgggagga	gagactgctg	gagacatggg	gaaactgaga	cagataattc	720
25	aaagaggagg	aattctgatc	acagaaagaa	aaaaagctga	cttttccaca	aaagacattt	780
	atcaggacct	caaccggttg	ttgaaaggca	aaaagggaga	gcagatgaat	agtgtctgat	840
	tgccagaaat	ggagaatcag	gttgacgttt	catcactgtc	tgcggtaatc	aagtttttag	900
	aactcttata	agatgattcc	aactttggac	agtttgaact	gactactttt	gacttcagcc	960
	agtatatgaa	attggatatt	gcagcagtc	gagcccttaa	cctttttcag	ggttctgttg	1020
30	aagataccac	tggctctcag	tctctggctg	ccttgctgaa	taagtgtaaa	acccctcaag	1080
	gacaaagact	tgtaaacacg	tggattaagc	agcctctcat	ggataagaac	agaatagagg	1140
	agagattgaa	tttagtggaa	gctttttaga	aagatgcaga	attgaggcag	actttacaag	1200
	aagattttact	tctgcgattc	ccagatctta	accgacttgc	caagaagtgt	caaagacaag	1260
	cagcaaaactt	acaagattgt	taccgactct	atcagggtat	aaatcaacta	cctaattgta	1320
35	tacaggctct	ggaaaaacat	gaaggaaaac	accagaaatt	attgttggca	gtttttgtga	1380
	ctcctcttac	tgatcttcgt	tctgacttct	ccaagtttca	ggaaatgata	gaaacaactt	1440
	tagatatgga	tcagggtgaa	aaccatgaat	tccttgtaaa	accttcattt	gatcctaact	1500
	tcagtgaatt	aagagaaata	atgaatgact	tggaaaagaa	gatgcagtc	acattataaa	1560
	gtgcagccag	agatottggc	ttggaccctg	gcaaacagat	taaactggat	tccagtgcac	1620
40	agtttggata	ttactttcgt	gtaacctgta	aggaagaaaa	agtccttcgt	aacaataaaa	1680
	acttttagtac	tgtagatata	cagaagaatg	gtgttaaatt	taccaacagc	aaattgactt	1740
	cttttaaatga	agagtatacc	aaaaataaaa	cagaatatga	agaagccag	gatgccattg	1800
	ttaaagaaat	tgtcaatatt	tcttcaggct	atgtagaacc	aatgcagaca	ctcaatgatg	1860
	tgttagctca	gctagatgct	gttgctcagct	ttgctcacgt	gtcaaatgga	gcacctgttc	1920
45	catatgtacg	accagccatt	ttggagaag	gacaaggaag	aattatatta	aaagcatcca	1980
	ggcatgcttg	tgttgaagtt	caagatgaaa	ttgcatttat	tcctaataac	gtatactttg	2040
	aaaaagataa	acagatgttc	cacatcatta	ctggcccca	tatgggaggt	aatcaacat	2100
	atattcgaca	aactgggggtg	atagtactca	tggcccaaat	tgggtgtttt	gtgccatgtg	2160
	agtcagcaga	agtgtccatt	gtggactgca	tcttagcccg	agtaggggct	ggtgacagtc	2220
50	aattgaaaag	agtctccacg	ttcatgttgc	aaatgttgg	aactgcttct	atcctcaggt	2280
	ctgcaaccaa	agattcatta	ataatcatag	atgaattggg	aagaggaact	tctacctacg	2340
	atggattttg	gttagcatgg	gctatatcag	aatacattgc	aacaaagatt	ggtgcttttt	2400
	gcatgtttgc	aaccattttt	catgaactta	ctgccttggc	caatcagata	ccaactgtta	2460
	ataatctata	tgtcacagca	ctcaccactg	aagagacctt	aactatgctt	tatcagggtga	2520
55	agaaaggtgt	ctgtgatcaa	agttttggga	ttcatgttgc	agagcttgc	aatttcccta	2580
	agcatgtaat	agagtgtgct	aaacagaaag	ccctggaaact	tgaggagttt	cagtatatatt	2640
	gagaatcgca	aggatatgat	atcatggaac	cagcagcaaa	gaagtgtat	ctggaaagag	2700
	agcaaggtga	aaaaattatt	caggagtctc	tgtccaaggt	gaaacaaatg	ccctttactg	2760
	aatgtcaga	agaaaacatc	acaataaagt	taaaacagct	aaaagctgaa	gtaatatgcaa	2820
60	agaataatag	ctttgtaaat	gaaatcattt	cacgaataaa	agttactacg	tgaaaaatcc	2880
	cagtaaatgga	atgaaggtaa	tattgataag	ctattgtctg	taatagtgtt	atattgtttt	2940
	atattaaccc	tttttccata	gtgttaactg	tcagtgccca	tggtgtatca	acttaataag	3000
	atatttagta	atattttact	ttgaggacat	tttcaaagat	ttttattttg	aaaaatgaga	3060
	gctgtaactg	aggactgttt	gcaattgaca	taggcaataa	taagtgtatg	gctgaatttt	3120
65	ataaataaaa	tcatgtagtt	tgtgg				3145

## MLH1 (human) (SEQ ID NO:13)

	MSFVAGVIRR	LDETVVNRIA	AGEVIQRPAN	AIKEMIENCL	DAKSTSIQVI	VKEGGLKLIQ	60
	IQDNGTGIRK	EDLDIVCERF	TTSKLQSFED	LASISTYGFR	GEALASISHV	AHVTITTKTA	120
5	DGKCAYRASY	SDGKLKAPPK	PCAGNQGTQI	TVEDLFYNIA	TRRKALKNPS	EEYKGILEVV	180
	GRYSVHNAGI	SFSVKKQGET	VADVRTLPA	STVDNIRSIF	GNAVSRELIE	IGCEDKTLAF	240
	KMNGYISNAN	YSVKKCIFLL	FINHRLVEST	SLRKAIETVY	AAYLPKNTHP	FLYLSLEISP	300
	QNVVDNVHPT	KHEVHFLHEE	SILERVQQHI	ESKLLGSNSS	RMFTQTLLP	GLAGPSGEMV	360
	KSTTSLTSSS	TSGSSDKVYA	HQMVRTDSRE	QKLDLFLQPL	SKPLSSQPOA	IVTEDKTDIS	420
10	SGRARQQDEE	MLELPAPAEV	AAKNQSLGEG	TTKGTSEMSE	KRGPTSSNPR	KRHREDSDE	480
	MVEDDSRKEM	TAACTPRRRI	INLTSVLSLQ	EEINEQGHEV	LREMLHNHSF	VGCVPNPQWAL	540
	AQHQTLYLL	NTTKLSEELF	YQILIYDFAN	FGVLRLESPA	PLFDLAMLAL	DSPESGWTEE	600
	DGPKEGLAEY	IVEFLKKKAE	MLADYFSLFI	DEEGNLIGLP	LLIDNYPVPL	EGLPIFILRL	660
	ATEVNWDEEK	ECFESLSKEC	AMFYISIRKQY	ISEESTLSGQ	QSEVPGSIPN	SWKWTVEHIV	720
15	YKALRSHILP	PKHFTEDGNI	LQLANLPDLY	KVFERC			756

## MLH1 (human) (SEQ ID NO:14)

	cttggctctt	ctggcgccaa	aatgtcggtc	gtggcagggg	ttattcgccg	gctggacgag	60
	acagtgggtga	accgcatcgc	ggcgggggaa	gttatccagc	ggccagctaa	tgctatcaaa	120
20	gagatgattg	agaactgttt	agatgcaaaa	tccacaagta	ttcaagtgat	tgtaaagag	180
	ggaggcctga	agttgattca	gatccaagac	aatggcaccg	ggatcaggaa	agaagatctg	240
	gatattgtat	gtgaaagggt	cactactagt	aaactgcagt	cctttgagga	tttagccagt	300
	atctctacac	atggctttcg	aggtgaggct	ttggccagca	taagccatgt	ggctcatgtt	360
	actattacaa	cgaaaacagc	tgatggaaaag	tgtgcataca	gagcaagtta	ctcagatgga	420
25	aaactgaaag	cccctcctaa	accatgtgct	ggcaatcaag	ggacccagat	caagggtggag	480
	gacotTTTTT	acaacatagc	cacgaggaga	aaagctttta	aaaatccaag	tgaagaatat	540
	gggaaaattt	tggaagttgt	tggcaggtat	tcagtacaca	atgcaggcat	tagtttctca	600
	gttaaaaaaac	aaggagagac	agtagctgat	gttaggacac	taccatgatc	ctcaaccgtg	660
	gacaatattc	gctccatctt	tggaatgtct	gttagtcgag	aactgataga	aattggatgt	720
30	gaggataaaa	ccctagcctt	caaaatgaat	ggttacatat	ccaatgcaaa	ctactcagtg	780
	aagaagtgc	tcttcttact	cttcatcaac	catcgtctgg	tagaatcaac	ttccttgaga	840
	aaagccatag	aaacagtgta	tgacgcctat	ttgccccaaa	acacacaccc	attcctgtac	900
	ctcagtttag	aaatcagtc	ccagaatgtg	gatgttaatg	tgacccccac	aaagcatgaa	960
	gttcaacttc	tgacagagga	gagcatcctg	gagcgggtgc	agcagcacat	cgagagcaag	1020
35	ctcctgggct	ccaattcctc	caggatgtac	ttcacccaga	ccttgctacc	taacttgctc	1080
	ggccccctctg	gggagatggt	taaatccaca	acaagtctga	cctcgtcttc	tacttctgga	1140
	agtagtgata	aggtctatgc	ccaccagatg	gttcgtacag	attcccggga	acagaagctt	1200
	gatgcatttc	tgacgcctct	gagcaaaccc	ctgtccagtc	agccccaggc	cattgtcaca	1260
	gaggataaga	cagatatttc	tagtggcagg	gctaggcagc	aagatgagga	gatgcttgaa	1320
40	ctcccagccc	ctgctgaagt	ggctgccaaa	aatcagagct	tgagggggga	tacaacaaag	1380
	gggagctcag	aaatgtcaga	gaagagagga	cctacttcca	gcaaccccag	aaagagacat	1440
	cgggaagatt	ctgatgtgga	aatggtggaa	gatgattccc	gaaaggaaat	gactgcagct	1500
	tgtaccccc	ggagaaggat	cattaacctc	actagtgttt	tgagtctcca	ggaagaaatt	1560
	aatgagcagg	gacatgaggt	tctcggggag	atgttgcata	accactcctt	cgtgggctgt	1620
45	gtgaatcctc	agtgggcctt	ggcacagcat	caaaccaagt	tataccttct	caacaccacc	1680
	aagcttagtg	agaactgtt	ctaccagata	ctcatttatg	atcttgccaa	ttttgggtgt	1740
	ctcaggttat	cggagccagc	accgtctttt	gacottgcca	tgcttgctct	agatagtgca	1800
	gagagtggct	ggacagagga	agatggtccc	aaagaaggac	ttgctgaata	cattgttgag	1860
	tttctgaaga	agaaggctga	gatgcttgca	gactatttct	ccttggaat	tgatgaggaa	1920
50	gggaacctga	ttggattacc	ccttctgatt	gacaactatg	tgcccccttt	ggagggactg	1980
	cctatottca	ttcttcgact	agccactgag	gtgaattggg	acgaagaaaa	ggaatgtttt	2040
	gaaagcctca	gtaaagaatg	cgctatgttc	tattccatcc	ggaagcagta	catatctgag	2100
	gagtcgaccc	tctcaggcca	gcagagtga	gtgcctggct	ccattccaaa	ctcctggaag	2160
	tggaactgtg	aacacattgt	ctataaagcc	ttgcgctcac	acattctgcc	tcctaaacat	2220
55	ttcacagaag	atggaaatat	cctgcagctt	gctaacctgc	ctgatctata	caaagtcttt	2280
	gagaggtgtt	aaatatggtt	atctatgcac	tgtgggatgt	gttcttcttt	ctctgtattc	2340
	cgatacaaag	tgttgatatca	aagtgtgata	tacaaagtgt	accaacataa	gtgttggtag	2400
	cacttaagac	ttatacttgc	cttctgatag	tattccttta	tacacagtgg	attgattata	2460
60	aataaataga	tgtgtcttaa	cata				2484

## hPMS2-134 (human) (SEQ ID NO:15)

```

MERAESSSTE PAKAIKPIDR KSVHQICSGQ VVLSLSTAVK ELVENS LDAG ATNIDLK LKD 60
YGVDLIEVSD NGCGVEEENF EGLTLKHHTS KIQEFADLTQ VETFGFRGEA LSSLCALSDV 120
TISTCHASAK VGT 133

```

## 5 hPMS2-134 (human cDNA) (SEQ ID NO:16)

```

cgaggcggat cgggtgttgc atccatggag cgagctgaga gctcgagtac agaacctgct 60
aaggccatca aacctattga tcggaagtca gtccatcaga tttgctctgg gcaggtggta 120
ctgagtctaa gcactgcggg aaaggagtta gtagaaaaca gtctggatgc tggtgccact 180
aatattgata taaagcttaa ggactatgga gtggatctta ttgaagtttc agacaatgga 240
10 tgtggggtag aagaagaaaa cttcgaaggc ttaactctga aacatcacac atctaagatt 300
caagagtttg ccgacctaac tcaggttgaa acttttggct ttcgggggga agctctgagc 360
tcactttgtg cactgagcga tgtcaccatt tctacctgcc acgcatcggc gaaggttgga 420
acttga 426

```

15 For further information on the background of the invention the following references may be consulted, each of which is incorporated herein by reference in its entirety:

1. Glaser, V. (1996) Can ReoPro repolish tarnished monoclonal therapeutics? *Nat. Biotechnol.* 14:1216-1217.
2. Weiner, L.M. (1999) Monoclonal antibody therapy of cancer. *Semin. Oncol.* 26:43-51.
3. Saez-Llorens, X.E. *et al.* (1998) Safety and pharmacokinetics of an intramuscular humanized monoclonal antibody to respiratory syncytial virus in premature infants and infants with bronchopulmonary dysplasia. *Pediat. Infect. Dis. J.* 17:787-791.
4. Shield, C.F. *et al.* (1996) A cost-effective analysis of OKT3 induction therapy in cadaveric kidney transplantation. *Am. J. Kidney Dis.* 27:855-864.
5. Khazaeli, M.B. *et al.* (1994) Human immune response to monoclonal antibodies. *J. Immunother.* 15:42-52.
6. Emery, S.C. and W.J. Harris "Strategies for humanizing antibodies" In: ANTIBODY ENGINEERING C.A.K. Borrebaeck (Ed.) Oxford University Press, N.Y. 1995, pp. 159-183.
7. U.S. Patent No. 5,530,101 to Queen and Selick.
8. Reff, M.E. (1993) High-level production of recombinant immunoglobulins in mammalian cells. *Curr. Opin. Biotechnol.* 4:573-576.
9. Neuberger, M. and M. Gruggermann, (1997) Monoclonal antibodies. Mice perform a human repertoire. *Nature* 386:25-26.

10. Fiedler, U. and U. Conrad (1995) High-level production and long-term storage of engineered antibodies in transgenic tobacco seeds. *Bio/Technology* 13:1090-1093.
11. Baker S.M. *et al.* (1995) Male defective in the DNA mismatch repair gene PMS2 exhibit abnormal chromosome synapsis in meiosis. *Cell* 82:309-319.
- 5 12. Bronner, C.E. *et al.* (1994) Mutation in the DNA mismatch repair gene homologue *hMLH1* is associated with hereditary non-polyposis colon cancer. *Nature* 368:258-261.
13. de Wind N. *et al.* (1995) Inactivation of the mouse *Msh2* gene results in mismatch repair deficiency, methylation tolerance, hyperrecombination, and predisposition to cancer. *Cell* 82:321-300.
- 10 14. Drummond, J.T. *et al.* (1995) Isolation of an hMSH2-p160 heterodimer that restores mismatch repair to tumor cells. *Science* 268:1909-1912.
15. Modrich, P. (1994) Mismatch repair, genetic stability, and cancer. *Science* 266:1959-1960.
- 15 16. Nicolaides, N.C. *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype. *Mol. Cell. Biol.* 18:1635-1641.
17. Prolla, T.A. *et al.* (1994) MLH1, PMS1, and MSH2 Interaction during the initiation of DNA mismatch repair in yeast. *Science* 264:1091-1093.
18. Strand, M. *et al.* (1993) Destabilization of tracts of simple repetitive DNA in yeast by mutations affecting DNA mismatch repair. *Nature* 365:274-276.
- 20 19. Su, S.S., R.S. Lahue, K.G. Au, and P. Modrich (1988) Mismatch specificity of methyl directed DNA mismatch corrections in vitro. *J. Biol. Chem.* 263:6829-6835.
20. Parsons, R. *et al.* (1993) Hypermutability and mismatch repair deficiency in RER<sup>+</sup> tumor cells. *Cell* 75:1227-1236.
- 25 21. Papadopoulos, N. *et al.* (1993) Mutation of a mutL homolog is associated with hereditary colon cancer. *Science* 263:1625-1629.
22. Perucho, M. (1996) Cancer of the microsatellite mutator phenotype. *Biol. Chem.* 377:675-684.
23. Nicolaides N.C., K.W. Kinzler, and B. Vogelstein (1995) Analysis of the 5' region of PMS2 reveals heterogenous transcripts and a novel overlapping gene. *Genomics* 29:329-334.
- 30 24. Nicolaides, N.C. *et al.* (1995) Genomic organization of the human PMS2 gene family. *Genomics* 30:195-206.

25. Palombo, F. *et al.* (1994) Mismatch repair and cancer. *Nature* 36:417.
26. Eshleman J.R. and S.D. Markowitz (1996) Mismatch repair defects in human carcinogenesis. *Hum. Mol. Genet.* 5:1489-494.
27. Liu, T. *et al.* (2000) Microsatellite instability as a predictor of a mutation in a DNA mismatch repair gene in familial colorectal cancer. *Genes Chromosomes Cancer* 27:17-25.
28. Nicolaides, N.C. *et al.* (1992) The Jun family members, c-JUN and JUND, transactivate the human *c-myc* promoter via an Ap1 like element. *J. Biol. Chem.* 267:19665-19672.
29. Shields, R.L. *et al.* (1995) Anti-IgE monoclonal antibodies that inhibit allergen-specific histamine release. *Int. Arch. Allergy Immunol.* 107:412-413.
30. Frigerio L. *et al.* (2000) Assembly, secretion, and vacuolar delivery of a hybrid immunoglobulin in plants. *Plant Physiol.* 123:1483-1494.
31. Bignami M, (2000) Unmasking a killer: DNA O(6)-methylguanine and the cytotoxicity of methylating agents. *Mutat. Res.* 462:71-82.
32. Drummond, J.T. *et al.* (1996) Cisplatin and adriamycin resistance are associated with MutLa and mismatch repair deficiency in an ovarian tumor cell line. *J. Biol. Chem.* 271:9645-19648.
33. Galio, L. *et al.* (1999) ATP hydrolysis-dependent formation of a dynamic ternary nucleoprotein complex with MutS and MutL. *Nucl. Acids Res.* 27:2325-23231.

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples which are provided herein for purposes of illustration only, and are not intended to limit the scope of the invention.

#### **EXAMPLE 1: Stable expression of dominant negative MMR genes in hybridoma cells**

It has been previously shown by Nicolaides *et al.* (Nicolaides *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype *Mol. Cell. Biol.* 18:1635-1641) that the expression of a dominant negative allele in an otherwise MMR proficient cell could render these host cells MMR deficient. The creation of MMR deficient cells can lead to the generation of genetic alterations

throughout the entire genome of a host organisms offspring, yielding a population of genetically altered offspring or siblings that may produce biochemicals with altered properties. This patent application teaches of the use of dominant negative MMR genes in antibody-producing cells, including but not limited to rodent hybridomas, human  
5 hybridomas, chimeric rodent cells producing human immunoglobulin gene products, human cells expressing immunoglobulin genes, mammalian cells producing single chain antibodies, and prokaryotic cells producing mammalian immunoglobulin genes or chimeric immunoglobulin molecules such as those contained within single-chain antibodies. The cell expression systems described above that are used to produce  
10 antibodies are well known by those skilled in the art of antibody therapeutics.

To demonstrate the ability to create MMR defective hybridomas using dominant negative alleles of MMR genes, we first transfected a mouse hybridoma cell line that is known to produce and antibody directed against the human IgE protein with an expression vector containing the human PMS2 (cell line referred to as HBPMS2), the previously  
15 published dominant negative PMS2 mutant referred herein as PMS134 (cell line referred to as HB134), or with no insert (cell line referred to as HBvec). The results showed that the PMS134 mutant could indeed exert a robust dominant negative effect, resulting in biochemical and genetic manifestations of MMR deficiency. Unexpectedly was the finding that the full length PMS2 also resulted in a lower MMR activity while no effect  
20 was seen in cells containing the empty vector. A brief description of the methods is provided below.

The MMR proficient mouse H36 hybridoma cell line was transfected with various *hPMS2* expression plasmids plus reporter constructs for assessing MMR activity. The MMR genes were cloned into the pEF expression vector, which contains  
25 the elongation factor promoter upstream of the cloning site followed by a mammalian polyadenylation signal. This vector also contains the NEO<sup>r</sup> gene that allows for selection of cells retaining this plasmid. Briefly, cells were transfected with 1 µg of each vector using polyliposomes following the manufacturer's protocol (Life Technologies). Cells were then selected in 0.5 mg/ml of G418 for 10 days and G418  
30 resistant cells were pooled together to analyze for gene expression. The pEF construct contains an intron that separates the exon 1 of the EF gene from exon 2, which is juxtaposed to the 5' end of the polylinker cloning site. This allows for a rapid reverse transcriptase polymerase chain reaction (RT-PCR) screen for cells expressing the spliced

products. At day 17, 100,000 cells were isolated and their RNA extracted using the trizol method as previously described (Nicolaidis N.C., Kinzler, K.W., and Vogelstein, B. (1995) Analysis of the 5' region of PMS2 reveals heterogeneous transcripts and a novel overlapping gene. *Genomics* 29:329-334). RNAs were reverse transcribed using

5 Superscript II (Life Technologies) and PCR amplified using a sense primer located in exon 1 of the EF gene (5'-ttt cgc aac ggg ttg gcc g-3') and an antisense primer (5'-gtt tca gag tta agc ctt cg-3') centered at nt 283 of the published human PMS2 cDNA, which will detect both the full length as well as the PMS134 gene expression. Reactions were carried out using buffers and conditions as previously described (Nicolaidis, N.C., *et*

10 *al.* (1995) Genomic organization of the human PMS2 gene family. *Genomics* 30:195-206), using the following amplification parameters: 94°C for 30 sec, 52°C for 2 min, 72°C for 2 min, for 30 cycles. Reactions were analyzed on agarose gels. Figure 1 shows a representative example of PMS expression in stably transduced H36 cells.

Expression of the protein encoded by these genes were confirmed via western

15 blot using a polyclonal antibody directed to the first 20 amino acids located in the N-terminus of the protein following the procedures previously described (data not shown) (Nicolaidis *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype. *Mol. Cell. Biol.* 18:1635-1641.

## 20 **EXAMPLE 2: hPMS134 Causes a Defect in MMR Activity and hypermutability in hybridoma cells**

A hallmark of MMR deficiency is the generation of unstable microsatellite repeats in the genome of host cells. This phenotype is referred to as microsatellite instability (MI) (Modrich, P. (1994) Mismatch repair, genetic stability, and cancer *Science*

25 266:1959-1960; Palombo, F., *et al.* (1994) Mismatch repair and cancer *Nature* 36:417). MI consists of deletions and/or insertions within repetitive mono-, di- and/or tri nucleotide repetitive sequences throughout the entire genome of a host cell. Extensive genetic analysis eukaryotic cells have found that the only biochemical defect that is capable of producing MI is defective MMR (Strand, M., *et al.* (1993) Destabilization of tracts of

30 simple repetitive DNA in yeast by mutations affecting DNA mismatch repair *Nature* 365:274-276; Perucho, M. (1996) Cancer of the microsatellite mutator phenotype. *Biol Chem.* 377:675-684; Eshleman J.R., and Markowitz, S.D. (1996) Mismatch repair defects in human carcinogenesis. *Hum. Mol. Genet.* 5:1489-494). In light of this unique feature



that defective MMR has on promoting MI, it is now used as a biochemical marker to survey for lack of MMR activity within host cells (Perucho, M. (1996) Cancer of the microsatellite mutator phenotype. *Biol Chem.* 377:675-684; Eshleman J.R., and Markowitz, S.D. (1996) Mismatch repair defects in human carcinogenesis. *Hum. Mol. Genet.* 5:1489-494; Liu, T., *et al.* (2000) Microsatellite instability as a predictor of a mutation in a DNA mismatch repair gene in familial colorectal cancer *Genes Chromosomes Cancer* 27:17-25).

A method used to detect MMR deficiency in eukaryotic cells is to employ a reporter gene that has a polynucleotide repeat inserted within the coding region that disrupts its reading frame due to a frame shift. In the case where MMR is defective, the reporter gene will acquire random mutations (i.e. insertions and/or deletions) within the polynucleotide repeat yielding clones that contain a reporter with an open reading frame. We have employed the use of an MMR-sensitive reporter gene to measure for MMR activity in HBvec, HBPMS2, and HBPMS134 cells. The reporter construct used the pCAR-OF, which contains a hygromycin resistance (HYG) gene plus a  $\beta$ -galactosidase gene containing a 29 bp out-of-frame poly-CA tract at the 5' end of its coding region. The pCAR-OF reporter would not generate  $\beta$ -galactosidase activity unless a frame-restoring mutation (i.e., insertion or deletion) arose following transfection. HBvec, HBPMS2, and HB134 cells were each transfected with pCAR-OF vector in duplicate reactions following the protocol described in Example 1. Cells were selected in 0.5 mg/ml G418 and 0.5mg/ml HYG to select for cells retaining both the MMR effector and the pCAR-OF reporter plasmids. All cultures transfected with the pCAR vector resulted in a similar number of HYG/G418 resistant cells. Cultures were then expanded and tested for  $\beta$ -galactosidase activity *in situ* as well as by biochemical analysis of cell extracts. For *in situ* analysis, 100,000 cells were harvested and fixed in 1% gluteraldehyde, washed in phosphate buffered saline solution and incubated in 1 ml of X-gal substrate solution [0.15 M NaCl, 1 mM MgCl<sub>2</sub>, 3.3 mM K<sub>4</sub>Fe(CN)<sub>6</sub>, 3.3 mM K<sub>3</sub>Fe(CN)<sub>6</sub>, 0.2% X-Gal ] in 24 well plates for 2 hours at 37°C. Reactions were stopped in 500 mM sodium bicarbonate solution and transferred to microscope slides for analysis. Three fields of 200 cells each were counted for blue ( $\beta$ -galactosidase positive cells) or white ( $\beta$ -galactosidase negative cells) to assess for MMR inactivation. Table 1 shows the results from these studies. While no  $\beta$ -galactosidase positive cells were observed in HBvec cells, 10% of the cells per

field were  $\beta$ -galactosidase positive in HB134 cultures and 2% of the cells per field were  $\beta$ -galactosidase positive in HBPMS2 cultures.

Cell extracts were prepared from the above cultures to measure  $\beta$ -galactosidase using a quantitative biochemical assay as previously described (Nicolaidis *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype *Mol. Cell. Biol.* 18:1635-1641; Nicolaidis, N.C., *et al.* (1992) The Jun family members, c-JUN and JUND, transactivate the human *c-myc* promoter via an Ap1 like element. *J. Biol. Chem.* 267:19665-19672). Briefly, 100,000 cells were collected, centrifuged and resuspended in 200  $\mu$ l of 0.25M Tris, pH 8.0. Cells were lysed by freeze/thawing three times and supernatants collected after microfugation at 14,000 rpms to remove cell debris. Protein content was determined by spectrophotometric analysis at OD<sup>280</sup>. For biochemical assays, 20  $\mu$ g of protein was added to buffer containing 45 mM 2-mercaptoethanol, 1mM MgCl<sub>2</sub>, 0.1 M NaPO<sub>4</sub> and 0.6 mg/ml Chlorophenol red- $\beta$ -D-galactopyranoside (CPRG, Boehringer Mannheim). Reactions were incubated for 1 hour, terminated by the addition of 0.5 M Na<sub>2</sub>CO<sub>3</sub>, and analyzed by spectrophotometry at 576 nm. H36 cell lysates were used to subtract out background. Figure 2 shows the  $\beta$ -galactosidase activity in extracts from the various cell lines. As shown, the HB134 cells produced the highest amount of  $\beta$ -galactosidase, while no activity was found in the HBvec cells containing the pCAR-OF. These data demonstrate the ability to generate MMR defective hybridoma cells using dominant negative MMR gene alleles.

**Table 1.**  $\beta$ -galactosidase expression of HBvec, HBPMS2 and HB134 cells transfected with pCAR-OF reporter vectors. Cells were transfected with the pCAR-OF  $\beta$ -galactosidase reporter plasmid. Transfected cells were selected in hygromycin and G418, expanded and stained with X-gal solution to measure for  $\beta$ -galactosidase activity (blue colored cells). 3 fields of 200 cells each were analyzed by microscopy. The results below represent the mean +/- standard deviation of these experiments.

produced by each clone. Reactions are stopped by adding 50  $\mu$ ls of 500mM sodium bicarbonate and analyzed by OD at 415nm using a BioRad plate reader. Clones exhibiting an enhanced signal over background cells (H36 control cells) are then isolated and expanded into 10 ml cultures for additional characterization and confirmation of ELISA data in triplicate experiments. ELISAs are also performed on conditioned (CM) from the same clones to measure total Ig production within the conditioned medium of each well. Clones that produce an increased ELISA signal and have increased antibody levels are then further analyzed for variants that over-express and/or over-secrete antibodies as described in Example 4. Analysis of five 96-well plates each from HBvec or HB134 cells have found that a significant number of clones with a higher Optical Density (OD) value is observed in the MMR-defective HB134 cells as compared to the HBvec controls. Figure 4 shows a representative example of HB134 clones producing antibodies that bind to specific antigen (in this case IgE) with a higher affinity. Figure 4 provides raw data from the analysis of 96 wells of HBvec (left graph) or HB134 (right graph) which shows 2 clones from the HB134 plate to have a higher OD reading due to 1) genetic alteration of the antibody variable domain that leads to an increased binding to IgE antigen, or 2) genetic alteration of a cell host that leads to over-production/secretion of the antibody molecule. Anti-Ig ELISA found that the two clones, shown in figure 4 have Ig levels within their CM similar to the surrounding wells exhibiting lower OD values. These data suggest that a genetic alteration occurred within the antigen binding domain of the antibody which in turn allows for higher binding to antigen.

Clones that produced higher OD values as determined by ELISA were further analyzed at the genetic level to confirm that mutations within the light or heavy chain variable region have occurred that lead to a higher binding affinity hence yielding to a stronger ELISA signal. Briefly, 100,000 cells are harvested and extracted for RNA using the Triazol method as described above. RNAs are reverse transcribed using Superscript II as suggested by the manufacturer (Life Technology) and PCR amplified for the antigen binding sites contained within the variable light and heavy chains. Because of the heterogeneous nature of these genes, the following degenerate primers are used to amplify light and heavy chain alleles from the parent H36 strain.

Light chain sense: 5'-GGA TTT TCA GGT GCA GAT TTT CAG-3' (SEQ ID NO:1)

Light chain antisense: 5'-ACT GGA TGG TGG GAA GAT GGA-3' (SEQ ID NO:2)

Heavy chain sense: 5'-A(G/T) GTN (A/C)AG CTN CAG (C/G)AG TC-3' (SEQ ID NO:3)

- 5 Heavy chain antisense: 5'-TNC CTT G(A/G)C CCC AGT A(G/A)(A/T)C-3' (SEQ ID NO:4)

- PCR reactions using degenerate oligonucleotides are carried out at 94°C for 30 sec, 52°C for 1 min, and 72°C for 1 min for 35 cycles. Products are analyzed on agarose gels.
- 10 Products of the expected molecular weights are purified from the gels by Gene Clean (Bio 101), cloned into T-tailed vectors, and sequenced to identify the wild type sequence of the variable light and heavy chains. Once the wild type sequence has been determined, non-degenerate primers were made for RT-PCR amplification of positive HB134 clones. Both the light and heavy chains were amplified, gel purified and sequenced using the
- 15 corresponding sense and antisense primers. The sequencing of RT-PCR products gives representative sequence data of the endogenous immunoglobulin gene and not due to PCR induced mutations. Sequences from clones were then compared to the wild type sequence for sequence comparison. An example of the ability to create *in vivo* mutations within an immunoglobulin light or heavy chain is shown in figure 5, where HB134 clone92 was
- 20 identified by ELISA to have an increased signal for hIgE. The light chain was amplified using specific sense and antisense primers. The light chain was RT-PCR amplified and the resulting product was purified and analyzed on an automated ABI377 sequencer. As shown in clone A, a residue -4 upstream of the CDR region 3 had a genetic change from ACT to TCT, which results in a Thr to Ser change within the framework region just
- 25 preceding the CDR#3. In clone B, a residue -6 upstream of the CDR region had a genetic change from CCC to CTC, which results in a Pro to His change within framework region preceding CDR#2.

- The ability to generate random mutations in immunoglobulin genes or chimeric immunoglobulin genes is not limited to hybridomas. Nicolaides et al. (Nicolaides *et al.*
- 30 (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype *Mol. Cell. Biol.* 18:1635-1641) has previously shown the ability to generate hypermutable hamster cells and produce mutations within an endogenous gene. A common method for producing humanized antibodies is to graft CDR sequences from a MAAb

(produced by immunizing a rodent host) onto a human Ig backbone, and transfection of the chimeric genes into Chinese Hamster Ovary (CHO) cells which in turn produce a functional Ab that is secreted by the CHO cells (Shields, R.L., *et al.* (1995) Anti-IgE monoclonal antibodies that inhibit allergen-specific histamine release. *Int. Arch. Allergy Immunol.*

5 107:412-413). The methods described within this application are also useful for generating genetic alterations within Ig genes or chimeric Igs transfected within host cells such as rodent cell lines, plants, yeast and prokaryotes (Frigerio L, *et al.* (2000) Assembly, secretion, and vacuolar delivery of a hybrid immunoglobulin in plants. *Plant Physiol.* 123:1483-1494).

10 These data demonstrate the ability to generate hypermutable hybridomas, or other Ig producing host cells that can be grown and selected, to identify structurally altered immunoglobulins yielding antibodies with enhanced biochemical properties, including but not limited to increased antigen binding affinity. Moreover, hypermutable clones that contain missense mutations within the immunoglobulin gene that result in an amino acid  
15 change or changes can be then further characterized for *in vivo* stability, antigen clearance, on-off binding to antigens, etc. Clones can also be further expanded for subsequent rounds of *in vivo* mutations and can be screened using the strategy listed above.

The use of chemical mutagens to produce genetic mutations in cells or whole organisms are limited due to the toxic effects that these agents have on "normal" cells. The  
20 use of chemical mutagens such as MNU in MMR defective organisms is much more tolerable yielding to a 10 to 100 fold increase in genetic mutation over MMR deficiency alone (Bignami M, (2000) Unmasking a killer: DNA O(6)-methylguanine and the cytotoxicity of methylating agents. *Mutat. Res.* 462:71-82). This strategy allows for the use of chemical mutagens to be used in MMR-defective Ab producing cells as a method for  
25 increasing additional mutations within immunoglobulin genes or chimeras that may yield functional Abs with altered biochemical properties such as enhanced binding affinity to antigen, etc.

**Example 4: Generation of antibody producing cells with enhanced antibody production**

Analysis of clones from H36 and HB134 following the screening strategy listed above has identified a significant number of clones that produce enhanced amounts of antibody into the medium. While a subset of these clones gave higher Ig binding data as determined by ELISA as a consequence of mutations within the antigen binding domains contained in the variable regions, others were found to contain "enhanced" antibody production. A summary of the clones producing enhanced amounts of secreted MAb is shown in TABLE 2, where a significant number of clones from HB134 cells were found to produce enhanced Ab production within the conditioned medium as compared to H36 control cells.

**TABLE 2.** Generation of hybridoma cells producing high levels of antibody. HB134 clones were assayed by ELISA for elevated Ig levels. Analysis of 480 clones showed that a significant number of clones had elevated MAb product levels in their CM. Quantification showed that several of these clones produced greater than 500ngs/ml of MAb due to either enhanced expression and/or secretion as compared to clones from the H36 cell line.

**Table 2. Production of MAb in CM from H36 and HB134 clones.**

Cell Line	% clones > 400 ng/ml	% clones >500 ng/ml
H36	1/480 = 0.2%	0/480 = 0%
HB134	50/480 = 10%	8/480 = 1.7%

20

Cellular analysis of HB134 clones with higher MAb levels within the conditioned medium (CM) were analyzed to determine if the increased production was simply due to genetic alterations at the Ig locus that may lead to over-expression of the polypeptides forming the antibody, or due to enhanced secretion due to a genetic alteration affecting secretory pathway mechanisms. To address this issue, we expanded three HB134 clones that had increased levels of antibody within their CM. 10,000 cells were prepared for western blot analysis to assay for intracellular steady state Ig protein levels (Figure 6). In addition, H36 cells were used as a standard reference (Lane 2) and a rodent fibroblast (Lane 1) was used as an Ig negative control. Briefly, cells were pelleted by centrifugation and lysed directly in 300  $\mu$ l of SDS lysis buffer (60 mM Tris, pH 6.8, 2% SDS, 10% glycerol,

30

0.1 M 2-mercaptoethanol, 0.001% bromophenol blue) and boiled for 5 minutes. Lysate proteins were separated by electrophoresis on 4-12% NuPAGE gels (for analysis of Ig heavy chain. Gels were electroblotted onto Immobilon-P (Millipore) in 48 mM Tris base, 40 mM glycine, 0.0375% SDS, 20% methanol and blocked at room temperature for 1 hour in Tris-buffered saline (TBS) plus 0.05% Tween-20 and 5% condensed milk. Filters were probed with a 1:10,000 dilution of sheep anti-mouse horseradish peroxidase conjugated monoclonal antibody in TBS buffer and detected by chemiluminescence using Supersignal substrate (Pierce). Experiments were repeated in duplicates to ensure reproducibility. Figure 6 shows a representative analysis where a subset of clones had enhanced Ig production which accounted for increased Ab production (Lane 5) while others had a similar steady state level as the control sample, yet had higher levels of Ab within the CM. These data suggest a mechanism whereby a subset of HB134 clones contained a genetic alteration that in turn produces elevated secretion of antibody.

The use of chemical mutagens to produce genetic mutations in cells or whole organisms are limited due to the toxic effects that these agents have on "normal" cells. The use of chemical mutagens such as MNU in MMR defective organisms is much more tolerable yielding to a 10 to 100 fold increase in genetic mutation over MMR deficiency alone (Bignami M, (2000) Unmasking a killer: DNA O(6)-methylguanine and the cytotoxicity of methylating agents. *Mutat. Res.* 462:71-82). This strategy allows for the use of chemical mutagens to be used in MMR-defective Ab producing cells as a method for increasing additional mutations within immunoglobulin genes or chimeras that may yield functional Abs with altered biochemical properties such as enhanced binding affinity to antigen, etc.

**Example 5: establishment of genetic stability in hybridoma cells with new output trait.**

The initial steps of MMR are dependent on two protein complexes, called MutS $\alpha$  and MutL $\alpha$  (Nicolaidis *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype. *Mol. Cell. Biol.* 18:1635-1641). Dominant negative MMR alleles are able to perturb the formation of these complexes with downstream biochemicals involved in the excision and polymerization of nucleotides comprising the "corrected" nucleotides. Examples from this application show the ability of a truncated MMR allele (PMS134) as well as a full length human PMS2 when expressed in

a hybridoma cell line is capable of blocking MMR resulting in a hypermutable cell line that gains genetic alterations throughout its entire genome per cell division. Once a cell line is produced that contains genetic alterations within genes encoding for an antibody, a single chain antibody, over expression of immunoglobulin genes and/or enhanced secretion of antibody, it is desirable to restore the genomic integrity of the cell host. This can be achieved by the use of inducible vectors whereby dominant negative MMR genes are cloned into such vectors, introduced into Ab producing cells and the cells are cultured in the presence of inducer molecules and/or conditions. Inducible vectors include but are not limited to chemical regulated promoters such as the steroid inducible MMTV, tetracycline regulated promoters, temperature sensitive MMR gene alleles, and temperature sensitive promoters.

The results described above lead to several conclusions. First, expression of hPMS2 and PMS134 results in an increase in microsatellite instability in hybridoma cells. That this elevated microsatellite instability is due to MMR deficiency was proven by evaluation of extracts from stably transduced cells. The expression of PMS134 results in a polar defect in MMR, which was only observed using heteroduplexes designed to test repair from the 5' direction (no significant defect in repair from the 3' direction was observed in the same extracts) (Nicolaides *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype. *Mol. Cell. Biol.* 18:1635-1641). Interestingly, cells deficient in hMLH1 also have a polar defect in MMR, but in this case preferentially affecting repair from the 3' direction (Drummond, J.T, *et al.* (1996) Cisplatin and adriamycin resistance are associated with MutLa and mismatch repair deficiency in an ovarian tumor cell line. *J. Biol. Chem.* 271:9645-19648). It is known from previous studies in both prokaryotes and eukaryotes that the separate enzymatic components mediate repair from the two different directions. Our results, in combination with those of Drummond *et al.* (Shields, R.L., *et al.* (1995) Anti-IgE monoclonal antibodies that inhibit allergen-specific histamine release. *Int. Arch Allergy Immunol.* 107:412-413), strongly suggest a model in which 5' repair is primarily dependent on hPMS2 while 3' repair is primarily dependent on hMLH1. It is easy to envision how the dimeric complex between PMS2 and MLH1 might set up this directionality. The combined results also demonstrate that a defect in directional MMR is sufficient to produce a MMR defective phenotype and suggests that any MMR gene allele is useful to produce genetically altered hybridoma cells, or a cell line that is producing Ig gene products. Moreover, the use of



such MMR alleles will be useful for generating genetically altered Ig polypeptides with altered biochemical properties as well as cell hosts that produce enhanced amounts of antibody molecules.

Another method that is taught in this application is that ANY method used to block MMR can be performed to generate hypermutability in an antibody-producing cell that can lead to genetically altered antibodies with enhanced biochemical features such as but not limited to increased antigen binding, enhanced pharmacokinetic profiles, etc. These processes can also be used to generate antibody producer cells that have increased Ig expression as shown in Example 4, figure 6 and/or increased antibody secretion as shown in Table 2.

In addition, we demonstrate the utility of blocking MMR in antibody producing cells to increase genetic alterations within Ig genes that may lead to altered biochemical features such as, but not limited to, increased antigen binding affinities (Figure 5A and 5B).

The blockade of MMR in such cells can be through the use of dominant negative MMR gene alleles from any species including bacteria, yeast, protozoa, insects, rodents, primates, mammalian cells, and man. Blockade of MMR can also be generated through the use of antisense RNA or deoxynucleotides directed to any of the genes involved in the MMR biochemical pathway. Blockade of MMR can be through the use of polypeptides that interfere with subunits of the MMR complex including but not limited to antibodies.

Finally, the blockade of MMR may be through the use chemicals such as but not limited to nonhydrolyzable ATP analogs, which have been shown to block MMR (Galio, L, *et al.* (1999) ATP hydrolysis-dependent formation of a dynamic ternary nucleoprotein complex with MutS and MutL. *Nucl. Acids Res.* 27:2325-23231).

## WE CLAIM:

1. A method for making a hypermutable, antibody producing cell, comprising introducing into a cell capable of producing antibodies a polynucleotide comprising a dominant negative allele of a mismatch repair gene.
2. The method of claim 1 wherein said polynucleotide is introduced by transfection of a suspension of cells *in vitro*.
3. The method of claim 1 wherein said mismatch repair gene is *PMS2*.
4. The method of claim 1 wherein said mismatch repair gene is human *PMS2*.
5. The method of claim 1 wherein said mismatch repair gene is *MLH1*.
6. The method of claim 1 wherein said mismatch repair gene is *PMS1*.
7. The method of claim 1 wherein said mismatch repair gene is *MSH2*.
8. The method of claim 1 wherein said mismatch repair gene is *MSH2*.
9. The method of claim 4 wherein said allele comprises a truncation mutation.
10. The method of claim 4 wherein said allele comprises a truncation mutation at codon 134.
11. The method of claim 10 wherein said truncation mutation is a thymidine at nucleotide 424 of wild-type *PMS2*.
12. The method of claim 1 wherein said polynucleotide is introduced into a fertilized egg of an animal.
13. The method of claim 12 wherein the fertilized egg is subsequently implanted into a pseudo-pregnant female whereby the fertilized egg develops into a mature transgenic animal.
14. The method of claim 12 wherein said mismatch repair gene is *PMS2*.
15. The method of claim 12 wherein said mismatch repair gene is human *PMS2*.
16. The method of claim 12 wherein said mismatch repair gene is human *MLH1*.
17. The method of claim 12 wherein said mismatch repair gene is human *PMS1*.
18. The method of claim 11 wherein said mismatch repair gene is a human *mutL* homolog.
19. The method of claim 15 wherein said allele comprises a truncation mutation.
20. The method of claim 15 wherein said allele comprises a truncation mutation at codon 134.
21. The method of claim 19 wherein said truncation mutation is a thymidine at nucleotide 424 of wild-type *PMS2*.

22. The method of claim 1 wherein said capability is due to the co-introduction of an immunoglobulin gene into said cell.
23. A homogeneous culture of hypermutable, mammalian cells wherein said cells comprise a dominant negative allele of a mismatch repair gene.
24. The culture of hypermutable, mammalian cells of claim 23 wherein the mismatch repair gene is *PMS2*.
25. The culture of hypermutable, mammalian cells of claim 24 wherein the mismatch repair gene is human *PMS2*.
26. The culture of hypermutable, mammalian cells of claim 23 wherein the mismatch repair gene is *MLH1*.
27. The culture of hypermutable, mammalian cells of claim 23 wherein the mismatch repair gene is *PMS1*.
28. The culture of hypermutable, mammalian cells of claim 23 wherein the mismatch repair gene is a human *mutL* homolog.
29. The culture of hypermutable, mammalian cells of claim 23 wherein the cells express a protein consisting of the first 133 amino acids of hPMS2.
30. A method for generating a mutation in a gene affecting antibody production in an antibody-producing cell comprising:
  - growing a said cell comprising said gene and a dominant negative allele of a mismatch repair gene; and
  - testing the cell to determine whether said gene of interest harbors a mutation.
31. The method of claim 30 wherein the step of testing comprises analyzing a nucleotide sequence of said gene.
32. The method of claim 30 wherein the step of testing comprises analyzing mRNA transcribed from said gene.
33. The method of claim 30 wherein the step of testing comprises analyzing a protein encoded by the gene of interest.
34. The method of claim 30 wherein the step of testing comprises analyzing the phenotype of said gene.
35. The method of claim 30 wherein the step of testing comprises analyzing the binding activity of an antibody.

36. A method wherein a mammalian cell is made MMR defective by the process of introducing a polynucleotide comprising an antisense oligonucleotide targeted against an allele of a mismatch repair gene into a mammalian cell, whereby the cell becomes hypermutable.
37. The method of claim 36 wherein the step of testing comprises analyzing a nucleotide sequence of said gene.
38. The method of claim 36 wherein the step of testing comprises analyzing mRNA transcribed from said gene.
39. The method of claim 36 wherein the step of testing comprises analyzing a protein encoded by said gene.
40. The method of claim 36 wherein the step of testing comprises analyzing the phenotype of said gene.
41. The method of claim 36 wherein the step of testing comprises analyzing the binding activity of an antibody.
42. A method for generating a mutation in a gene affecting antibody production in an antibody-producing cell comprising:
  - growing said cell comprising said gene and a polynucleotide encoding a dominant negative allele of a mismatch repair gene; and
  - testing said cell to determine whether said cell harbors at least one mutation in said gene yielding to a new biochemical feature to the product of said gene, wherein said new biochemical feature is selected from the group consisting of over-expression of said product, enhanced secretion of said product, enhanced affinity of said product for antigen, and combinations thereof.
43. The method of claim 42 wherein the step of testing comprises analyzing the steady state expression of the immunoglobulin gene of said cell.
44. The method of claim 42 wherein the step of testing comprises analyzing steady state mRNA transcribed from the immunoglobulin gene of said cell.
45. The method of claim 42 wherein the step of testing comprises analyzing the amount of secreted protein encoded by the immunoglobulin gene of said cell.
46. The method of claim 36 wherein the cell is made by the process of introducing a polynucleotide comprising a dominant negative allele of a mismatch repair gene into a cell in the presence of DNA mutagens.

47. The method of claim 46 wherein the step of testing comprises analyzing a nucleotide sequence of an immunoglobulin gene of said cell.
48. The method of claim 46 wherein the step of testing comprises analyzing mRNA transcribed from the immunoglobulin gene of said cell.
49. The method of claim 46 wherein the step of testing comprises analyzing the immunoglobulin protein encoded by said gene.
50. The method of claim 46 wherein the step of testing comprises analyzing the biochemical activity of the protein encoded by said gene.
51. A hypermutable transgenic mammalian cell made by the method of claim 42.
52. The transgenic mammalian cell of claim 51 wherein said cell is from primate.
53. The transgenic mammalian cell of claim 51 wherein said cell is from rodent.
54. The transgenic mammalian cell of claim 51 wherein said cell is from human.
55. The transgenic mammalian cell of claim 51 wherein said cell is eukaryotic.
56. The transgenic mammalian cell of claim 51 wherein said cell is prokaryotic.
57. A method of reversibly altering the hypermutability of an antibody producing cell comprising introducing an inducible vector into a cell, wherein said inducible vector comprises a dominant negative allele of a mismatch repair gene operably linked to an inducible promoter, and inducing said cell to express said dominant negative mismatch repair gene.
58. The method of claim 57 wherein said mismatch repair gene is *PMS2*.
59. The method of claim 58 wherein said mismatch repair gene is human *PMS2*.
60. The method of claim 57 wherein said mismatch repair gene is *MLH1*.
61. The method of claim 57 wherein said mismatch repair gene is *PMS1*.
62. The method of claim 57 wherein said mismatch repair gene is a human *mutL* homolog.
63. The method of claim 57 wherein said cell expresses a protein consisting of the first 133 amino acids of hPMS2.
64. The method of claim 57 further comprising analyzing the immunoglobulin protein expressed by said antibody producing cell.
65. The method of claim 64 further comprising ceasing induction of said cell, thereby restoring genetic stability of said cell.
66. A method of producing genetically altered antibodies comprising

transfecting a polynucleotide encoding an immunoglobulin protein into a cell, wherein said cell comprises a dominant negative mismatch repair gene;

growing said cell, thereby producing a hypermutated polynucleotide encoding a hypermutated immunoglobulin protein;

screening for a desirable property of said hypermutated immunoglobulin protein;

isolating said hypermutated polynucleotide; and

transfecting said hypermutated polynucleotide into a genetically stable cell, thereby producing a hypermutated antibody-producing, genetically stable cell.

67. The method of claim 66 wherein said mismatch repair gene is *PMS2*.
68. The method of claim 66 wherein said mismatch repair gene is human *PMS2*.
69. The method of claim 66 wherein said mismatch repair gene is *MLH1*.
70. The method of claim 66 wherein said mismatch repair gene is *PMS1*.
71. The method of claim 66 wherein said mismatch repair gene is a human *mutL* homolog.
72. The method of claim 66 wherein said cell expresses a protein consisting of the first 133 amino acids of hPMS2.




FIG. 1






Fig. 2



Fig. 3



**Fig.4**



\* = clones with a significant difference in antigen binding





Fig. 5A

**Fig. 5B**




Fig. 6

## SEQUENCE LISTING

<110> Nicolaides, Nicholas C  
Grasso, Luigi  
Sass, Philip M

<120> METHODS FOR GENERATING GENETICALLY ALTERED  
ANTIBODY-PRODUCING CELL LINES WITH IMPROVED ANTIBODY  
CHARACTERISTICS

<130> MOR0004

<140> 00/000,000

<141> 2000-11-07

<160> 16

<170> PatentIn Ver. 2.1

<210> 1

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide  
primer

<400> 1

ggatttttcag gtgcagattt tcag

24

<210> 2

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide  
primer

<400> 2

actggatggt gggaagatgg a

21

<210> 3

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide  
primer

<400> 3

akgtnmagct ncagsagtc

19

<210> 4

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide  
primer

<400> 4

tnccttgrcc ccagtarwc

19

<210> 5

<211> 859

<212> PRT

<213> Mus musculus

<400> 5

Met Glu Gln Thr Glu Gly Val Ser Thr Glu Cys Ala Lys Ala Ile Lys  
1 5 10 15

Pro Ile Asp Gly Lys Ser Val His Gln Ile Cys Ser Gly Gln Val Ile  
20 25 30

Leu Ser Leu Ser Thr Ala Val Lys Glu Leu Ile Glu Asn Ser Val Asp  
35 40 45

Ala Gly Ala Thr Thr Ile Asp Leu Arg Leu Lys Asp Tyr Gly Val Asp  
50 55 60

Leu Ile Glu Val Ser Asp Asn Gly Cys Gly Val Glu Glu Glu Asn Phe  
65 70 75 80

Glu Gly Leu Ala Leu Lys His His Thr Ser Lys Ile Gln Glu Phe Ala  
85 90 95

Asp Leu Thr Gln Val Glu Thr Phe Gly Phe Arg Gly Glu Ala Leu Ser

100	105	110
Ser Leu Cys Ala Leu Ser Asp Val Thr Ile Ser Thr Cys His Gly Ser		
115	120	125
Ala Ser Val Gly Thr Arg Leu Val Phe Asp His Asn Gly Lys Ile Thr		
130	135	140
Gln Lys Thr Pro Tyr Pro Arg Pro Lys Gly Thr Thr Val Ser Val Gln		
145	150	155
His Leu Phe Tyr Thr Leu Pro Val Arg Tyr Lys Glu Phe Gln Arg Asn		
165	170	175
Ile Lys Lys Glu Tyr Ser Lys Met Val Gln Val Leu Gln Ala Tyr Cys		
180	185	190
Ile Ile Ser Ala Gly Val Arg Val Ser Cys Thr Asn Gln Leu Gly Gln		
195	200	205
Gly Lys Arg His Ala Val Val Cys Thr Ser Gly Thr Ser Gly Met Lys		
210	215	220
Glu Asn Ile Gly Ser Val Phe Gly Gln Lys Gln Leu Gln Ser Leu Ile		
225	230	235
Pro Phe Val Gln Leu Pro Pro Ser Asp Ala Val Cys Glu Glu Tyr Gly		
245	250	255
Leu Ser Thr Ser Gly Arg His Lys Thr Phe Ser Thr Phe Arg Ala Ser		
260	265	270
Phe His Ser Ala Arg Thr Ala Pro Gly Gly Val Gln Gln Thr Gly Ser		
275	280	285
Phe Ser Ser Ser Ile Arg Gly Pro Val Thr Gln Gln Arg Ser Leu Ser		
290	295	300
Leu Ser Met Arg Phe Tyr His Met Tyr Asn Arg His Gln Tyr Pro Phe		
305	310	315
Val Val Leu Asn Val Ser Val Asp Ser Glu Cys Val Asp Ile Asn Val		
325	330	335
Thr Pro Asp Lys Arg Gln Ile Leu Leu Gln Glu Glu Lys Leu Leu Leu		
340	345	350
Ala Val Leu Lys Thr Ser Leu Ile Gly Met Phe Asp Ser Asp Ala Asn		



355		360		365											
Lys	Leu	Asn	Val	Asn	Gln	Gln	Pro	Leu	Leu	Asp	Val	Glu	Gly	Asn	Leu
370		375		380											
Val	Lys	Leu	His	Thr	Ala	Glu	Leu	Glu	Lys	Pro	Val	Pro	Gly	Lys	Gln
385		390		395		400									
Asp	Asn	Ser	Pro	Ser	Leu	Lys	Ser	Thr	Ala	Asp	Glu	Lys	Arg	Val	Ala
		405		410		415									
Ser	Ile	Ser	Arg	Leu	Arg	Glu	Ala	Phe	Ser	Leu	His	Pro	Thr	Lys	Glu
		420		425		430									
Ile	Lys	Ser	Arg	Gly	Pro	Glu	Thr	Ala	Glu	Leu	Thr	Arg	Ser	Phe	Pro
	435			440		445									
Ser	Glu	Lys	Arg	Gly	Val	Leu	Ser	Ser	Tyr	Pro	Ser	Asp	Val	Ile	Ser
	450			455		460									
Tyr	Arg	Gly	Leu	Arg	Gly	Ser	Gln	Asp	Lys	Leu	Val	Ser	Pro	Thr	Asp
465		470		475		480									
Ser	Pro	Gly	Asp	Cys	Met	Asp	Arg	Glu	Lys	Ile	Glu	Lys	Asp	Ser	Gly
		485		490		495									
Leu	Ser	Ser	Thr	Ser	Ala	Gly	Ser	Glu	Glu	Glu	Phe	Ser	Thr	Pro	Glu
		500		505		510									
Val	Ala	Ser	Ser	Phe	Ser	Ser	Asp	Tyr	Asn	Val	Ser	Ser	Leu	Glu	Asp
	515			520		525									
Arg	Pro	Ser	Gln	Glu	Thr	Ile	Asn	Cys	Gly	Asp	Leu	Asp	Cys	Arg	Pro
	530			535		540									
Pro	Gly	Thr	Gly	Gln	Ser	Leu	Lys	Pro	Glu	Asp	His	Gly	Tyr	Gln	Cys
545		550		555		560									
Lys	Ala	Leu	Pro	Leu	Ala	Arg	Leu	Ser	Pro	Thr	Asn	Ala	Lys	Arg	Phe
		565		570		575									
Lys	Thr	Glu	Glu	Arg	Pro	Ser	Asn	Val	Asn	Ile	Ser	Gln	Arg	Leu	Pro
		580		585		590									
Gly	Pro	Gln	Ser	Thr	Ser	Ala	Ala	Glu	Val	Asp	Val	Ala	Ile	Lys	Met
	595			600		605									
Asn	Lys	Arg	Ile	Val	Leu	Leu	Glu	Phe	Ser	Leu	Ser	Ser	Leu	Ala	Lys

610	615	620
Arg Met Lys Gln Leu Gln His Leu Lys Ala Gln Asn Lys His Glu Leu		
625	630	635 640
Ser Tyr Arg Lys Phe Arg Ala Lys Ile Cys Pro Gly Glu Asn Gln Ala		
	645	650 655
Ala Glu Asp Glu Leu Arg Lys Glu Ile Ser Lys Ser Met Phe Ala Glu		
	660	665 670
Met Glu Ile Leu Gly Gln Phe Asn Leu Gly Phe Ile Val Thr Lys Leu		
	675	680 685
Lys Glu Asp Leu Phe Leu Val Asp Gln His Ala Ala Asp Glu Lys Tyr		
	690	695 700
Asn Phe Glu Met Leu Gln Gln His Thr Val Leu Gln Ala Gln Arg Leu		
	705 710	715 720
Ile Thr Pro Gln Thr Leu Asn Leu Thr Ala Val Asn Glu Ala Val Leu		
	725	730 735
Ile Glu Asn Leu Glu Ile Phe Arg Lys Asn Gly Phe Asp Phe Val Ile		
	740	745 750
Asp Glu Asp Ala Pro Val Thr Glu Arg Ala Lys Leu Ile Ser Leu Pro		
	755	760 765
Thr Ser Lys Asn Trp Thr Phe Gly Pro Gln Asp Ile Asp Glu Leu Ile		
	770	775 780
Phe Met Leu Ser Asp Ser Pro Gly Val Met Cys Arg Pro Ser Arg Val		
	785 790	795 800
Arg Gln Met Phe Ala Ser Arg Ala Cys Arg Lys Ser Val Met Ile Gly		
	805	810 815
Thr Ala Leu Asn Ala Ser Glu Met Lys Lys Leu Ile Thr His Met Gly		
	820	825 830
Glu Met Asp His Pro Trp Asn Cys Pro His Gly Arg Pro Thr Met Arg		
	835	840 845
His Val Ala Asn Leu Asp Val Ile Ser Gln Asn		
	850	855

<210> 6  
 <211> 3056  
 <212> DNA  
 <213> Mus musculus

<400> 6  
 gaattccggt gaaggtcctg aagaatttcc agattcctga gtatcattgg aggagacaga 60  
 taacctgtcg tcaggtaacg atgggtgtata tgcaacagaa atgggtgttc ctggagacgc 120  
 gtcttttccc gagagcggca ccgcaactct cccgcggtga ctgtgactgg aggagtcctg 180  
 catccatgga gcaaaccgaa ggcgtgagta cagaatgtgc taaggccatc aagcctattg 240  
 atgggaagtc agtccatcaa atttgttctg ggcaggtgat actcagttta agcaccgctg 300  
 tgaaggagtt gatagaaaat agtgtagatg ctggtgctac tactattgat ctaaggctta 360  
 aagactatgg ggtggacctc attgaagttt cagacaatgg atgtggggtg gaagaagaaa 420  
 actttgaagg tctagctctg aaacatcaca catctaagat tcaagagttt gccgacctca 480  
 cgcaggttga aactttcggc tttcgggggg aagctctgag ctctctgtgt gcactaagtg 540  
 atgtcactat atctacctgc cacgggtctg caagcgttgg gactcgactg gtgtttgacc 600  
 ataattgggaa aatcacccag aaaactccct acccccgacc taaaggaaac acagtcagtg 660  
 tgcagcactt attttataca ctaccctgct gttacaaaga gtttcagagg aacattaata 720  
 aggagtattc caaaatggtg caggtcttac aggcgtactg tatcatctca gcaggcgtcc 780  
 gtgtaagctg cactaatcag ctcgacaggg ggaagcggca cgtgtggtg tgcacaagcg 840  
 gcaogtcttg catgaaggaa aatatcgggt ctgtgtttgg ccagaagcag ttgcaaagcc 900  
 tcattccttt tgttcagctg cccctagtgt acgctgtgtg tgaagagtao gccctgagca 960  
 cttcaggacg ccacaaaacc tttctacgt ttcgggcttc atttcacagt gcacgcacgg 1020  
 cgccgggagg agtgcaacag acaggcagtt tttcttcac aatcagaggc cctgtgacct 1080  
 agcaaaggtc tctaagcttg tcaatgaggt tttatcacat gtataaccgg catcagtacc 1140  
 catttgctgt ccttaacgtt tccgttgact cagaatgtgt ggatattaat gtaactccag 1200  
 ataaaaggca aattctacta caagaagaga agctattgct ggccgtttta aagacctcct 1260  
 tgataggaat gtttgacagt gatgcaaaca agcttaatgt caaccagcag cactgtctag 1320  
 atgttgaagg taacttagta aagctgcata ctgcagaact agaaaagcct gtgccaggaa 1380  
 agcaagataa ctctccttca ctgaagagca cagcagacga gaaaagggtg gcatccatct 1440  
 ccaggctgag agaggccttt tctcttcac ctactaaaga gatcaagtct aggggtccag 1500  
 agactgctga actgacacgg agttttccaa gtgagaaaag gggcgtgtta tctcttatc 1560  
 cttcagacgt catctcttac agaggcctcc gtggctcgca ggacaaattg gtgagtccca 1620  
 cggacagccc tgggtactgt atggacagag agaaaataga aaaagactca gggctcagca 1680  
 gcacctcagc tggctctgag gaagagttca gcaccccgaga agtggccagt agcttttagca 1740  
 gtgactataa cgtgagctcc ctagaagaca gaccttotca ggaaaccata aactgtggtg 1800  
 acctggactg ccgtcctcca ggtacaggac agtccttgaa gccagaagac catggatatc 1860  
 aatgcaaagc tctacctcta gctcgtctgt caccacaaaa tgccaagcgc ttcaagacag 1920  
 aggaaagacc ctcaaatgtc aacattttctc aaagattgcc tggtcctcag agcacctcag 1980  
 cagctgaggt cgatgtagcc ataaaaatga ataagagaat cgtgctcctc gagttctctc 2040  
 tgagttctct agctaagcga atgaagcagt tacagcacct aaaggcgcag aacaaacatg 2100  
 aactgagtta cagaaaattt agggccaaga tttgccctgg agaaaaacca gcagcagaag 2160  
 atgaactcag aaaagagatt agtaaatcga tgtttgcaga gatggagatc ttgggtcagt 2220  
 ttaacctggg atttatagta accaaactga aagaggacct ctctctggtg gaccagcatg 2280  
 ctgcggatga gaagtacaac tttgagatgc tgcagcagca cacggtgctc caggcgcaga 2340  
 ggctcatcac accccagact ctgaacttaa ctgctgtcaa tgaagctgta ctgatagaaa 2400  
 atctggaaat attcagaaaag aatggctttg actttgtcat tgatgaggat gctccagtca 2460  
 ctgaaagggc taaattgatt tccttaccac ctagtaaaaa ctggaccttt ggacccaag 2520

atatagatga actgatcttt atgttaagtg acagccctgg ggtcatgtgc cggccctcac 2580  
 gagtcagaca gatgtttgct tccagagcct gtcggaagtc agtgatgatt ggaacggcgc 2640  
 tcaatgcgag cgagatgaag aagctcatca cccacatggg tgagatggac cacccttgga 2700  
 actgccccca cggcaggcca accatgaggc acgttgccaa tctggatgtc atctctcaga 2760  
 actgacacac ccctttagc atagagttta ttacagattg ttcggtttgc aaagagaagg 2820  
 ttttaagtaa tctgattatc gttgtacaaa aattagcatg ctgctttaat gtactggatc 2880  
 catttaaaag cagtgttaag gcaggcatga tggagtgttc ctctagctca gctacttggg 2940  
 tgatccggtg ggagctcatg tgagcccagg actttgagac cactccgagc cacattcatg 3000  
 agactcaatt caaggacaaa aaaaaaaga tattttttgaa gcctttttaa aaaaaa 3056

<210> 7

<211> 862

<212> PRT

<213> Homo sapiens

<400> 7

Met Glu Arg Ala Glu Ser Ser Ser Thr Glu Pro Ala Lys Ala Ile Lys  
 1 5 10 15

Pro Ile Asp Arg Lys Ser Val His Gln Ile Cys Ser Gly Gln Val Val  
 20 25 30

Leu Ser Leu Ser Thr Ala Val Lys Glu Leu Val Glu Asn Ser Leu Asp  
 35 40 45

Ala Gly Ala Thr Asn Ile Asp Leu Lys Leu Lys Asp Tyr Gly Val Asp  
 50 55 60

Leu Ile Glu Val Ser Asp Asn Gly Cys Gly Val Glu Glu Glu Asn Phe  
 65 70 75 80

Glu Gly Leu Thr Leu Lys His His Thr Ser Lys Ile Gln Glu Phe Ala  
 85 90 95

Asp Leu Thr Gln Val Glu Thr Phe Gly Phe Arg Gly Glu Ala Leu Ser  
 100 105 110

Ser Leu Cys Ala Leu Ser Asp Val Thr Ile Ser Thr Cys His Ala Ser  
 115 120 125

Ala Lys Val Gly Thr Arg Leu Met Phe Asp His Asn Gly Lys Ile Ile  
 130 135 140

Gln Lys Thr Pro Tyr Pro Arg Pro Arg Gly Thr Thr Val Ser Val Gln  
 145 150 155 160

Gln Leu Phe Ser Thr Leu Pro Val Arg His Lys Glu Phe Gln Arg Asn

8

420	425	430
Lys Pro His Ser Pro Lys Thr Pro Glu Pro Arg Arg Ser Pro Leu Gly		
435	440	445
Gln Lys Arg Gly Met Leu Ser Ser Ser Thr Ser Gly Ala Ile Ser Asp		
450	455	460
Lys Gly Val Leu Arg Pro Gln Lys Glu Ala Val Ser Ser Ser His Gly		
465	470	475 480
Pro Ser Asp Pro Thr Asp Arg Ala Glu Val Glu Lys Asp Ser Gly His		
	485	490 495
Gly Ser Thr Ser Val Asp Ser Glu Gly Phe Ser Ile Pro Asp Thr Gly		
	500	505 510
Ser His Cys Ser Ser Glu Tyr Ala Ala Ser Ser Pro Gly Asp Arg Gly		
	515	520 525
Ser Gln Glu His Val Asp Ser Gln Glu Lys Ala Pro Glu Thr Asp Asp		
	530	535 540
Ser Phe Ser Asp Val Asp Cys His Ser Asn Gln Glu Asp Thr Gly Cys		
	545	550 555 560
Lys Phe Arg Val Leu Pro Gln Pro Thr Asn Leu Ala Thr Pro Asn Thr		
	565	570 575
Lys Arg Phe Lys Lys Glu Glu Ile Leu Ser Ser Ser Asp Ile Cys Gln		
	580	585 590
Lys Leu Val Asn Thr Gln Asp Met Ser Ala Ser Gln Val Asp Val Ala		
	595	600 605
Val Lys Ile Asn Lys Lys Val Val Pro Leu Asp Phe Ser Met Ser Ser		
	610	615 620
Leu Ala Lys Arg Ile Lys Gln Leu His His Glu Ala Gln Gln Ser Glu		
	625	630 635 640
Gly Glu Gln Asn Tyr Arg Lys Phe Arg Ala Lys Ile Cys Pro Gly Glu		
	645	650 655
Asn Gln Ala Ala Glu Asp Glu Leu Arg Lys Glu Ile Ser Lys Thr Met		
	660	665 670
Phe Ala Glu Met Glu Ile Ile Gly Gln Phe Asn Leu Gly Phe Ile Ile		

675	680	685
Thr Lys Leu Asn Glu Asp Ile Phe Ile Val Asp Gln His Ala Thr Asp		
690	695	700
Glu Lys Tyr Asn Phe Glu Met Leu Gln Gln His Thr Val Leu Gln Gly		
705	710	715
Gln Arg Leu Ile Ala Pro Gln Thr Leu Asn Leu Thr Ala Val Asn Glu		
	725	730
Ala Val Leu Ile Glu Asn Leu Glu Ile Phe Arg Lys Asn Gly Phe Asp		
	740	745
Phe Val Ile Asp Glu Asn Ala Pro Val Thr Glu Arg Ala Lys Leu Ile		
	755	760
Ser Leu Pro Thr Ser Lys Asn Trp Thr Phe Gly Pro Gln Asp Val Asp		
	770	775
Glu Leu Ile Phe Met Leu Ser Asp Ser Pro Gly Val Met Cys Arg Pro		
	785	790
Ser Arg Val Lys Gln Met Phe Ala Ser Arg Ala Cys Arg Lys Ser Val		
	805	810
Met Ile Gly Thr Ala Leu Asn Thr Ser Glu Met Lys Lys Leu Ile Thr		
	820	825
His Met Gly Glu Met Asp His Pro Trp Asn Cys Pro His Gly Arg Pro		
	835	840
Thr Met Arg His Ile Ala Asn Leu Gly Val Ile Ser Gln Asn		
	850	855
		860

&lt;210&gt; 8

&lt;211&gt; 2771

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 8

```

cgaggcggat cgggtgttgc atccatggag cgagctgaga gctcgagtac agaacctgct 60
aaggccatca aacctattga tcggaagtca gtccatcaga ttgctctgg gcagggtgga 120
ctgagtctaa gcactgcggt aaaggagtta gtagaaaaca gtctggatgc tggtgccact 180
aatattgatc taaagcttaa ggactatgga gtggatctta ttgaagtctc agacaatgga 240
tgtggggtag aagaagaaaa cttcgaaggc ttaactctga aacatcacac atctaagatt 300
caagagtttg ccgacctaac tcaggttgaa acttttggct ttcgggggga agctctgagc 360

```

```

tcactttgtg cactgagcga tgtcaccatt tctacctgcc acgcacggc gaaggttga 420
actcgactga tgtttgatca caatgggaaa attatccaga aaaccccta ccccgcccc 480
agagggacca cagtcagcgt gcagcagtta ttttccacac tacctgtgcg ccataaggaa 540
tttcaaagga atattaagaa ggagtatgcc aaaatggtcc aggtcttaca tgcatactgt 600
atcatttcag caggcatccg tgtaagttgc accaatcagc ttggacaagg aaaacgacag 660
cctgtggtat gcacaggtgg aagccccagc ataaaggaaa atatcggtc tgtgtttggg 720
cagaagcagt tgcaaagcct cattcctttt gttcagctgc cccctagtga ctccgtgtgt 780
gaagagtacg gtttgagctg ttcggtatgt ctgcataatc ttttttacat ctccaggtttc 840
atttcacaat gcacgcagtg agttggaagg agttcaacag acagacagtt tttctttatc 900
aaccggcggc cttgtgacct agcaaaggtc tgcagactcg tgaatgaggt ctaccacatg 960
tataatcgac accagtatcc atttgttgtt cttaacatct ctgttgattc agaatgcgtt 1020
gatatcaatg ttactccaga taaaaggcaa attttgctac aagaggaaaa gctttttgtt 1080
gcagttttta agacctcttt gataggaatg tttgatagt atgtcaacaa gctaaatgtc 1140
agtcagcagc cactgctgga tgttgaagg aacttaataa aaatgcatgc agcggatttt 1200
gaaaagccca tggtagaaaa gcaggatcaa tccccctcat taaggactgg agaagaaaaa 1260
aaagacgtgt ccatttccag actgcgagag gccttttctc ttcgtcacac aacagagaac 1320
aagcctcaca gcccaaagac tccagaacca agaaggagcc ctctaggaca gaaaaggggt 1380
atgctgtctt ctagcacttc aggtgccatc tctgacaaag gcgtcctgag acctcagaaa 1440
gaggcagtga gttccagtca cggaccagc gacctacgg acagagcggg ggtggagaag 1500
gactcggggc acggcagcac ttccgtggat tctgaggggt tcagcatccc agacacgggc 1560
agtcactgca gcagcgagta tgcggccagc tccccagggg acaggggctc gcaggaacat 1620
gtggactctc aggagaaagc gcctgaaact gacgactct tttcagatgt ggaactgcat 1680
tcaaaccagg aagataccgg atgtaaatct cgagttttgc ctccagccac taatctcgca 1740
accccaaaaca caaagcgttt taaaaaagaa gaaattcttt ccagttctga cttttgtcaa 1800
aagttagtaa atactcagga catgtcagcc tctcaggttg atgtagctgt gaaaattaat 1860
aagaaagttg tgcccctgga cttttctatg agttctttag ctaaacgaat aaagcagtta 1920
catcatgaag cacagcaaag tgaaggggaa cagaattaca ggaagtttag ggcaâagatt 1980
tgtcctggag aaaatcaagc agccgaagat gaactaagaa aagagataag taaaacgatg 2040
tttgcgaaaa tggaaatcat tggtcagttt aacctgggat ttataataac caaactgaat 2100
gaggatatct tcatagtgga ccagcatgcc acggacgaga agtataactt cgagatgctg 2160
cagcagcaca ccgtgctcca ggggcagagg ctcatagcac ctccagactct caacttaact 2220
gctgttaatg aagctgttct gatagaaat ctggaaatat ttagaaagaa tggctttgat 2280
tttgttatcg atgaaatgc tccagtcact gaaagggcta aactgatttc cttgccaact 2340
agtaaaaact ggaccttcgg accccaggac gtcgatgaac tgatcttcat gctgagcgac 2400
agccctgggg tcatgtgccg gccttccga gtcaagcaga tgtttgcctc cagagcctgc 2460
cggaagtggg tgatgattgg gactgctctt aacacaagcg agatgaagaa actgatcacc 2520
cacatggggg agatggacca cccctggaac tgtcccatg gaaggccaac catgagacac 2580
atcgccaacc tgggtgtcat ttctcagaac tgaccgtagt cactgtatgg aataattgg 2640
tttatcgagc atttttatgt tttgaaagac agagtcttca ctaacctttt ttgttttaaa 2700
atgaaacctg ctacttaaaa aaaatacaca tcacacccat ttaaaagtga tottgagaac 2760
cttttcaaac c

```

```

<210> 9      0
<211> 932
<212> PRT
<213> Homo sapiens

```



&lt;400&gt; 9

```

Met Lys Gln Leu Pro Ala Ala Thr Val Arg Leu Leu Ser Ser Ser Gln
 1             5             10             15

Ile Ile Thr Ser Val Val Ser Val Val Lys Glu Leu Ile Glu Asn Ser
      20             25             30

Leu Asp Ala Gly Ala Thr Ser Val Asp Val Lys Leu Glu Asn Tyr Gly
      35             40             45

Phe Asp Lys Ile Glu Val Arg Asp Asn Gly Glu Gly Ile Lys Ala Val
      50             55             60

Asp Ala Pro Val Met Ala Met Lys Tyr Tyr Thr Ser Lys Ile Asn Ser
      65             70             75             80

His Glu Asp Leu Glu Asn Leu Thr Thr Tyr Gly Phe Arg Gly Glu Ala
      85             90             95

Leu Gly Ser Ile Cys Cys Ile Ala Glu Val Leu Ile Thr Thr Arg Thr
      100            105            110

Ala Ala Asp Asn Phe Ser Thr Gln Tyr Val Leu Asp Gly Ser Gly His
      115            120            125

Ile Leu Ser Gln Lys Pro Ser His Leu Gly Gln Gly Thr Thr Val Thr
      130            135            140

Ala Leu Arg Leu Phe Lys Asn Leu Pro Val Arg Lys Gln Phe Tyr Ser
      145            150            155            160

Thr Ala Lys Lys Cys Lys Asp Glu Ile Lys Lys Ile Gln Asp Leu Leu
      165            170            175

Met Ser Phe Gly Ile Leu Lys Pro Asp Leu Arg Ile Val Phe Val His
      180            185            190

Asn Lys Ala Val Ile Trp Gln Lys Ser Arg Val Ser Asp His Lys Met
      195            200            205

Ala Leu Met Ser Val Leu Gly Thr Ala Val Met Asn Asn Met Glu Ser
      210            215            220

Phe Gln Tyr His Ser Glu Glu Ser Gln Ile Tyr Leu Ser Gly Phe Leu
      225            230            235            240

Pro Lys Cys Asp Ala Asp His Ser Phe Thr Ser Leu Ser Thr Pro Glu
      245            250            255

```

Arg	Ser	Phe	Ile	Phe	Ile	Asn	Ser	Arg	Pro	Val	His	Gln	Lys	Asp	Ile	260	265	270
Leu	Lys	Leu	Ile	Arg	His	His	Tyr	Asn	Leu	Lys	Cys	Leu	Lys	Glu	Ser	275	280	285
Thr	Arg	Leu	Tyr	Pro	Val	Phe	Phe	Leu	Lys	Ile	Asp	Val	Pro	Thr	Ala	290	295	300
Asp	Val	Asp	Val	Asn	Leu	Thr	Pro	Asp	Lys	Ser	Gln	Val	Leu	Leu	Gln	305	310	315
Asn	Lys	Glu	Ser	Val	Leu	Ile	Ala	Leu	Glu	Asn	Leu	Met	Thr	Thr	Cys	325	330	335
Tyr	Gly	Pro	Leu	Pro	Ser	Thr	Asn	Ser	Tyr	Glu	Asn	Asn	Lys	Thr	Asp	340	345	350
Val	Ser	Ala	Ala	Asp	Ile	Val	Leu	Ser	Lys	Thr	Ala	Glu	Thr	Asp	Val	355	360	365
Leu	Phe	Asn	Lys	Val	Glu	Ser	Ser	Gly	Lys	Asn	Tyr	Ser	Asn	Val	Asp	370	375	380
Thr	Ser	Val	Ile	Pro	Phe	Gln	Asn	Asp	Met	His	Asn	Asp	Glu	Ser	Gly	385	390	395
Lys	Asn	Thr	Asp	Asp	Cys	Leu	Asn	His	Gln	Ile	Ser	Ile	Gly	Asp	Phe	405	410	415
Gly	Tyr	Gly	His	Cys	Ser	Ser	Glu	Ile	Ser	Asn	Ile	Asp	Lys	Asn	Thr	420	425	430
Lys	Asn	Ala	Phe	Gln	Asp	Ile	Ser	Met	Ser	Asn	Val	Ser	Trp	Glu	Asn	435	440	445
Ser	Gln	Thr	Glu	Tyr	Ser	Lys	Thr	Cys	Phe	Ile	Ser	Ser	Val	Lys	His	450	455	460
Thr	Gln	Ser	Glu	Asn	Gly	Asn	Lys	Asp	His	Ile	Asp	Glu	Ser	Gly	Glu	465	470	475
Asn	Glu	Glu	Glu	Ala	Gly	Leu	Glu	Asn	Ser	Ser	Glu	Ile	Ser	Ala	Asp	485	490	495
Glu	Trp	Ser	Arg	Gly	Asn	Ile	Leu	Lys	Asn	Ser	Val	Gly	Glu	Asn	Ile	500	505	510

Glu Pro Val Lys Ile Leu Val Pro Glu Lys Ser Leu Pro Cys Lys Val  
 515 520 525

Ser Asn Asn Asn Tyr Pro Ile Pro Glu Gln Met Asn Leu Asn Glu Asp  
 530 535 540

Ser Cys Asn Lys Lys Ser Asn Val Ile Asp Asn Lys Ser Gly Lys Val  
 545 550 555 560

Thr Ala Tyr Asp Leu Leu Ser Asn Arg Val Ile Lys Lys Pro Met Ser  
 565 570 575

Ala Ser Ala Leu Phe Val Gln Asp His Arg Pro Gln Phe Leu Ile Glu  
 580 585 590

Asn Pro Lys Thr Ser Leu Glu Asp Ala Thr Leu Gln Ile Glu Glu Leu  
 595 600 605

Trp Lys Thr Leu Ser Glu Glu Glu Lys Leu Lys Tyr Glu Glu Lys Ala  
 610 615 620

Thr Lys Asp Leu Glu Arg Tyr Asn Ser Gln Met Lys Arg Ala Ile Glu  
 625 630 635 640

Gln Glu Ser Gln Met Ser Leu Lys Asp Gly Arg Lys Lys Ile Lys Pro  
 645 650 655

Thr Ser Ala Trp Asn Leu Ala Gln Lys His Lys Leu Lys Thr Ser Leu  
 660 665 670

Ser Asn Gln Pro Lys Leu Asp Glu Leu Leu Gln Ser Gln Ile Glu Lys  
 675 680 685

Arg Arg Ser Gln Asn Ile Lys Met Val Gln Ile Pro Phe Ser Met Lys  
 690 695 700

Asn Leu Lys Ile Asn Phe Lys Lys Gln Asn Lys Val Asp Leu Glu Glu  
 705 710 715 720

Lys Asp Glu Pro Cys Leu Ile His Asn Leu Arg Phe Pro Asp Ala Trp  
 725 730 735

Leu Met Thr Ser Lys Thr Glu Val Met Leu Leu Asn Pro Tyr Arg Val  
 740 745 750

Glu Glu Ala Leu Leu Phe Lys Arg Leu Leu Glu Asn His Lys Leu Pro  
 755 760 765

Ala Glu Pro Leu Glu Lys Pro Ile Met Leu Thr Glu Ser Leu Phe Asn  
 770 775 780

Gly Ser His Tyr Leu Asp Val Leu Tyr Lys Met Thr Ala Asp Asp Gln  
 785 790 795 800

Arg Tyr Ser Gly Ser Thr Tyr Leu Ser Asp Pro Arg Leu Thr Ala Asn  
 805 810 815

Gly Phe Lys Ile Lys Leu Ile Pro Gly Val Ser Ile Thr Glu Asn Tyr  
 820 825 830

Leu Glu Ile Glu Gly Met Ala Asn Cys Leu Pro Phe Tyr Gly Val Ala  
 835 840 845

Asp Leu Lys Glu Ile Leu Asn Ala Ile Leu Asn Arg Asn Ala Lys Glu  
 850 855 860

Val Tyr Glu Cys Arg Pro Arg Lys Val Ile Ser Tyr Leu Glu Gly Glu  
 865 870 875 880

Ala Val Arg Leu Ser Arg Gln Leu Pro Met Tyr Leu Ser Lys Glu Asp  
 885 890 895

Ile Gln Asp Ile Ile Tyr Arg Met Lys His Gln Phe Gly Asn Glu Ile  
 900 905 910

Lys Glu Cys Val His Gly Arg Pro Phe Phe His His Leu Thr Tyr Leu  
 915 920 925

Pro Glu Thr Thr  
 930

&lt;210&gt; 10

&lt;211&gt; 3063

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 10

ggcacgagtg gctgcttgcg gctagtgat ggtaattgcc tgccctcgcg tagcagcaag 60  
 ctgctctgtt aaaagcgaaa atgaaacaat tgccctcgcg aacagttcga ctccctttcaa 120  
 gttctcagat catcacttcg gtggctcagtg ttgtaaaaga gcttattgaa aactccttgg 180  
 atgctggtgc cacaagcgta gatgttaaac tggagaacta tggatttgat aaaattgagg 240  
 tgcgagataa cggggagggt atcaaggctg ttgatgcacc tgtaatggca atgaagtact 300  
 acacotcaaa aataaatagt catgaagatc ttgaaaattt gacaacttac ggttttcgtg 360  
 gagaagcctt ggggtcaatt tggtgtatag ctgaggtttt aattacaaca agaacggctg 420

```

ctgataat ttt tagcaccag tatgttttag atggcagtgg ccacatactt tctcagaaac 480
cttcacatct tgggtcaaggt acaactgtaa ctgctttaag attattttaag aatctacctg 540
taagaaagca gttttactca actgcaaaaa aatgtaaaga tgaaataaaa aagatccaag 600
atctcctcat gagctttggt atccttaaac ctgacttaag gattgtcttt gtacataaca 660
aggcagttat ttggcagaaa agcagagtat cagatcaca gatggctctc atgtcagttc 720
tggggactgc tgttatgaac aatatggaat cctttcagta ccactctgaa gaatctcaga 780
tttatctcag tggatttctt ccaaagtgtg atgcagacca ctctttcact agtctttcaa 840
caccagaaaag aagtttcatc ttcataaaca gtgcaccagt acatcaaaaa gatattctta 900
agttaatccg acatcattac aatctgaaat gcctaaaagga atctactcgt ttgtatcctg 960
ttttctttct gaaaatcgat gtccctacag ctgatgttga tgtaaattta acaccagata 1020
aaagccaagt attattacaa aataaggaat ctgttttaat tgctcttgaa aatctgatga 1080
cgacttgta tggaccatta cctagtacaa attcttatga aaataataaa acagatgttt 1140
ccgcagctga catcgttctt agtaaaacag cagaaacaga tgtgcttttt aataaagtgg 1200
aatcatctgg aaagaattat tcaaatgttg atacttcagt cattccattc caaaatgata 1260
tgcataatga tgaatctgga aaaaacactg atgattgttt aaatcaccag ataagtattg 1320
gtgactttgg ttatggctcat tgtagttagt aaatttctaa cattgataaa aacactaaga 1380
atgcatttca ggacatttca atgagtaatg tatcatggga gaactctcag acggaatata 1440
gtaaaacttg ttttataagt tccgttaagc acaccagtc agaaaatggc aataaagacc 1500
atatagatga gagtggggaa aatgaggaag aagcaggtct tgaaaactct tcggaaattt 1560
ctgcagatga gtggagcagg ggaaatatac ttaaaaattc agtgggagag aatattgaac 1620
ctgtgaaaat tttagtgcct gaaaaaagtt taccatgtaa agtaagtaat aataattatc 1680
caatccctga acaaatgaat cttaatgaag attcatgtaa caaaaaatca aatgtaatag 1740
ataataaatc tggaaaagtt acagcttatg atttacttag caatcgagta atcaagaaac 1800
ccatgtcagc aagtgtctct tttgttcaag atcatcgtcc tcagtttctc atagaaaatc 1860
ctaagactag tttagaggat gcaacactac aaattgaaga actgtggaag acattgagtg 1920
aagaggaaaa actgaaatat gaagagaagg ctactaaaga cttggaacga tacaatagtc 1980
aaatgaagag agccattgaa caggagtcac aaatgtcact aaaagatggc agaaaaaga 2040
taaaaccac cagcgcattg aatttggtcc agaagcaca gttaaaaacc tcattatcta 2100
atcaaccaa acttgatgaa ctcttcagt ccaaattga aaaaagaagg agtcaaaata 2160
ttaaaatggt acagatcccc tttcttatga aaaacttaaa aataaatttt aagaacaaa 2220
acaaagtga cttagaagag aaggatgaac cttgcttgat ccacaatctc aggtttcctg 2280
atgcatggct aatgacatcc aaaacagagg taatgttatt aaatccatat agagtagaag 2340
aagccctgct atttaaaaga cttcttgaga atcataaact tcctgcagag cactggaaa 2400
agccaattat gttaacagag agtcttttta atggatctca ttatttagac gttttatata 2460
aaatgacagc agatgaccaa agatacagtg gatcaactta cctgtctgat cctcgtctta 2520
cagcgaatgg tttcaagata aaattgatac caggagtttc aattactgaa aattacttgg 2580
aaatagaagg aatggctaat tgtctcccat tctatggagt agcagattta aaagaaattc 2640
ttaatgctat attaaacaga aatgcaaagg aagtttatga atgtagacct cgcaaagtga 2700
taagttat t agaggagaa gcagtgcgtc tatccagaca attacccatg tacttatcaa 2760
aagaggacat ccaagacatt atctacagaa tgaagcacca gtttggaat gaaattaaag 2820
agtgtgttca tggctgcccc ttttttcatc atttaaccta tcttcagaa actacatgat 2880
taaataatgt taagaagatt agttaccatt gaaattgggt ctgtcataaa acagcatgag 2940
tctgttttta aattatcttt gtattatgtg tcacatgggt attttttaaa tgaggattca 3000
ctgacttggt tttatattga aaaaagttcc acgtattgta gaaaacgtaa ataaactaat 3060
aac 3063

```

&lt;211&gt; 934

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 11

Met Ala Val Gln Pro Lys Glu Thr Leu Gln Leu Glu Ser Ala Ala Glu  
 1 5 10 15

Val Gly Phe Val Arg Phe Phe Gln Gly Met Pro Glu Lys Pro Thr Thr  
 20 25 30

Thr Val Arg Leu Phe Asp Arg Gly Asp Phe Tyr Thr Ala His Gly Glu  
 35 40 45

Asp Ala Leu Leu Ala Ala Arg Glu Val Phe Lys Thr Gln Gly Val Ile  
 50 55 60

Lys Tyr Met Gly Pro Ala Gly Ala Lys Asn Leu Gln Ser Val Val Leu  
 65 70 75 80

Ser Lys Met Asn Phe Glu Ser Phe Val Lys Asp Leu Leu Leu Val Arg  
 85 90 95

Gln Tyr Arg Val Glu Val Tyr Lys Asn Arg Ala Gly Asn Lys Ala Ser  
 100 105 110

Lys Glu Asn Asp Trp Tyr Leu Ala Tyr Lys Ala Ser Pro Gly Asn Leu  
 115 120 125

Ser Gln Phe Glu Asp Ile Leu Phe Gly Asn Asn Asp Met Ser Ala Ser  
 130 135 140

Ile Gly Val Val Gly Val Lys Met Ser Ala Val Asp Gly Gln Arg Gln  
 145 150 155 160

Val Gly Val Gly Tyr Val Asp Ser Ile Gln Arg Lys Leu Gly Leu Cys  
 165 170 175

Glu Phe Pro Asp Asn Asp Gln Phe Ser Asn Leu Glu Ala Leu Leu Ile  
 180 185 190

Gln Ile Gly Pro Lys Glu Cys Val Leu Pro Gly Gly Glu Thr Ala Gly  
 195 200 205

Asp Met Gly Lys Leu Arg Gln Ile Ile Gln Arg Gly Gly Ile Leu Ile  
 210 215 220

Thr Glu Arg Lys Lys Ala Asp Phe Ser Thr Lys Asp Ile Tyr Gln Asp

225		230		235		240
Leu Asn Arg Leu Leu Lys Gly Lys Lys Gly Glu Gln Met Asn Ser Ala						
	245		250			255
Val Leu Pro Glu Met Glu Asn Gln Val Ala Val Ser Ser Leu Ser Ala						
	260		265			270
Val Ile Lys Phe Leu Glu Leu Leu Ser Asp Asp Ser Asn Phe Gly Gln						
	275		280			285
Phe Glu Leu Thr Thr Phe Asp Phe Ser Gln Tyr Met Lys Leu Asp Ile						
	290		295			300
Ala Ala Val Arg Ala Leu Asn Leu Phe Gln Gly Ser Val Glu Asp Thr						
305		310		315		320
Thr Gly Ser Gln Ser Leu Ala Ala Leu Leu Asn Lys Cys Lys Thr Pro						
	325		330			335
Gln Gly Gln Arg Leu Val Asn Gln Trp Ile Lys Gln Pro Leu Met Asp						
	340		345			350
Lys Asn Arg Ile Glu Glu Arg Leu Asn Leu Val Glu Ala Phe Val Glu						
	355		360			365
Asp Ala Glu Leu Arg Gln Thr Leu Gln Glu Asp Leu Leu Arg Arg Phe						
	370		375			380
Pro Asp Leu Asn Arg Leu Ala Lys Lys Phe Gln Arg Gln Ala Ala Asn						
385		390		395		400
Leu Gln Asp Cys Tyr Arg Leu Tyr Gln Gly Ile Asn Gln Leu Pro Asn						
	405		410			415
Val Ile Gln Ala Leu Glu Lys His Glu Gly Lys His Gln Lys Leu Leu						
	420		425			430
Leu Ala Val Phe Val Thr Pro Leu Thr Asp Leu Arg Ser Asp Phe Ser						
	435		440			445
Lys Phe Gln Glu Met Ile Glu Thr Thr Leu Asp Met Asp Gln Val Glu						
	450		455			460
Asn His Glu Phe Leu Val Lys Pro Ser Phe Asp Pro Asn Leu Ser Glu						
465		470		475		480
Leu Arg Glu Ile Met Asn Asp Leu Glu Lys Lys Met Gln Ser Thr Leu						

	485		490		495														
Ile	Ser	Ala	Ala	Arg	Asp	Leu	Gly	Leu	Asp	Pro	Gly	Lys	Gln	Ile	Lys				
		500						505					510						
Leu	Asp	Ser	Ser	Ala	Gln	Phe	Gly	Tyr	Tyr	Phe	Arg	Val	Thr	Cys	Lys				
		515					520					525							
Glu	Glu	Lys	Val	Leu	Arg	Asn	Asn	Lys	Asn	Phe	Ser	Thr	Val	Asp	Ile				
		530				535				540									
Gln	Lys	Asn	Gly	Val	Lys	Phe	Thr	Asn	Ser	Lys	Leu	Thr	Ser	Leu	Asn				
545					550				555						560				
Glu	Glu	Tyr	Thr	Lys	Asn	Lys	Thr	Glu	Tyr	Glu	Glu	Ala	Gln	Asp	Ala				
				565				570						575					
Ile	Val	Lys	Glu	Ile	Val	Asn	Ile	Ser	Ser	Gly	Tyr	Val	Glu	Pro	Met				
		580					585						590						
Gln	Thr	Leu	Asn	Asp	Val	Leu	Ala	Gln	Leu	Asp	Ala	Val	Val	Ser	Phe				
		595				600						605							
Ala	His	Val	Ser	Asn	Gly	Ala	Pro	Val	Pro	Tyr	Val	Arg	Pro	Ala	Ile				
		610				615						620							
Leu	Glu	Lys	Gly	Gln	Gly	Arg	Ile	Ile	Leu	Lys	Ala	Ser	Arg	His	Ala				
625					630				635						640				
Cys	Val	Glu	Val	Gln	Asp	Glu	Ile	Ala	Phe	Ile	Pro	Asn	Asp	Val	Tyr				
				645				650						655					
Phe	Glu	Lys	Asp	Lys	Gln	Met	Phe	His	Ile	Ile	Thr	Gly	Pro	Asn	Met				
		660					665						670						
Gly	Gly	Lys	Ser	Thr	Tyr	Ile	Arg	Gln	Thr	Gly	Val	Ile	Val	Leu	Met				
		675				680						685							
Ala	Gln	Ile	Gly	Cys	Phe	Val	Pro	Cys	Glu	Ser	Ala	Glu	Val	Ser	Ile				
		690				695					700								
Val	Asp	Cys	Ile	Leu	Ala	Arg	Val	Gly	Ala	Gly	Asp	Ser	Gln	Leu	Lys				
705					710				715						720				
Gly	Val	Ser	Thr	Phe	Met	Ala	Glu	Met	Leu	Glu	Thr	Ala	Ser	Ile	Leu				
				725				730						735					
Arg	Ser	Ala	Thr	Lys	Asp	Ser	Leu	Ile	Ile	Ile	Asp	Glu	Leu	Gly	Arg				



740                                      745                                      750  
 Gly Thr Ser Thr Tyr Asp Gly Phe Gly Leu Ala Trp Ala Ile Ser Glu  
           755                                      760                                      765  
 Tyr Ile Ala Thr Lys Ile Gly Ala Phe Cys Met Phe Ala Thr His Phe  
           770                                      775                                      780  
 His Glu Leu Thr Ala Leu Ala Asn Gln Ile Pro Thr Val Asn Asn Leu  
 785                                      790                                      795                                      800  
 His Val Thr Ala Leu Thr Thr Glu Glu Thr Leu Thr Met Leu Tyr Gln  
                                     805                                      810                                      815  
 Val Lys Lys Gly Val Cys Asp Gln Ser Phe Gly Ile His Val Ala Glu  
                                     820                                      825                                      830  
 Leu Ala Asn Phe Pro Lys His Val Ile Glu Cys Ala Lys Gln Lys Ala  
                                     835                                      840                                      845  
 Leu Glu Leu Glu Glu Phe Gln Tyr Ile Gly Glu Ser Gln Gly Tyr Asp  
                                     850                                      855                                      860  
 Ile Met Glu Pro Ala Ala Lys Lys Cys Tyr Leu Glu Arg Glu Gln Gly  
 865                                      870                                      875                                      880  
 Glu Lys Ile Ile Gln Glu Phe Leu Ser Lys Val Lys Gln Met Pro Phe  
                                     885                                      890                                      895  
 Thr Glu Met Ser Glu Glu Asn Ile Thr Ile Lys Leu Lys Gln Leu Lys  
                                     900                                      905                                      910  
 Ala Glu Val Ile Ala Lys Asn Asn Ser Phe Val Asn Glu Ile Ile Ser  
                                     915                                      920                                      925  
 Arg Ile Lys Val Thr Thr  
                                     930

&lt;210&gt; 12

&lt;211&gt; 3145

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 12

ggcggggaaac agcttagtggt gtgtgggggtc ggcgattttc ttcaaccagg aggtgaggag 60  
 gtttcgacat ggcggtgcag ccgaaggaga cgctgcagtt ggagagcgcg gccgaggtcg 120  
 gcttcgtgcg cttctttcag ggcattgccgg agaagccgac caccacagtg cgccttttcg 180

```

accggggcga cttctatacg ggcacggcg aggcgcgct gctggccgcc cgggaggtgt 240
tcaagaccca gggggtgacg aagtacatgg ggccggcagg agcaaagaat ctgcagagtg 300
ttgtgcttag taaaatgaat tttgaatcct ttgtaaaaga tcttctctctg gttcgtcagt 360
atagagttga agtttataag aatagagctg gaaataaggg atccaaggag aatgattggg 420
atttggcata taaggcttct cctggcaatc tctctcagtt tgaagacatt ctctttggta 480
acaatgatat gtcagcttcc attggtgttg tgggtgttaa aatgtccgca gttgatggcc 540
agagacagggt tggagttggg tatgtggatt ccatacagag gaaactagga ctgtgtgaat 600
tccctgataa tgatcagttc tccaatcttg aggcctcct catccagatt ggaccaaagg 660
aatgtgtttt acccgaggga gagactgctg gagacatggg gaaactgaga cagataattc 720
aaagaggagg aattctgacg acagaaagaa aaaaagctga cttttccaca aaagacattt 780
atcaggacct caaccggttg ttgaaaggca aaaagggaga gcagatgaat agtgctgtat 840
tgccagaaat ggagaatcag gttgcagttt catcactgtc tgcggtaatc aagtttttag 900
aactcttata agatgattcc aactttggac agtttgaact gactactttt gacttcagcc 960
agtatatgaa attggatatt gcagcagtc gagcccttaa cttttttcag ggttctgttg 1020
aagataccac tggctctcag tctctggctg ccttgctgaa taagtgtaaa acccctcaag 1080
gacaaagact tgtaaccag tggattaagc agcctctcat ggataagaac agaataagg 1140
agagattgaa tttagtggaa gctttttag aagatgcaga attgaggcag actttacaag 1200
aagatttact tcgtcgattc ccagatctta accgacttgc caagaagttt caaagacaag 1260
cagcaaactt acaagattgt taccgactct atcagggtat aaatcaacta cctaattgta 1320
tacaggctct ggaaaaacat gaaggaaaac accagaaatt attgttggca gtttttgtga 1380
ctcctottac tgatcttcgt tctgacttct ccaagtttca ggaaatgata gaaacaactt 1440
tagatatgga tcaggtggaa aaccatgaat tccttgtaaa accttcattt gatcctaata 1500
tcagtgaatt aagagaaata atgaatgact tggaaaagaa gatgcagtc acattaataa 1560
gtgcagccag agatcttggc ttggaccctg gcaaacagat taaactggat tccagtgcac 1620
agtttggata ttactttcgt gtaacctgta aggaagaaaa agtccttcgt aacaataaaa 1680
acttttagtac tgtagatata cagaagaatg gtgttaaatt taccaacagc aaattgactt 1740
ctttaaatga agagtatacc aaaaataaaa cagaatatga agaagcccag gatgccattg 1800
ttaaagaaat tgtcaatatt tcttcaggct atgtagaacc aatgcagaca ctaaatgatg 1860
tgtagctca gctagatgct gttgtcagct ttgctcacgt gtcaaatgga gcacctgttc 1920
catatgtacg accagccatt ttggagaaag gacaagggaag aattatatta aaagcatcca 1980
ggcatgcttg tgttgaagtt caagatgaaa ttgcatttat tctaatagac gtatactttg 2040
aaaaagataa acagatgttc cacatcatta ctggccccaa tatgggaggt aaatcaacat 2100
atattcgaca aactggggtg atagtaactc tggcccaaatt tgggtgtttt gtgccatgtg 2160
agtcagcaga agtgtccatt gtggactgca tcttagcccg agtaggggct ggtgacagtc 2220
aattgaaagg agtctccacg ttcattggctg aaatgttga aactgcttct atcctcagg 2280
ctgcaaccaa agattcatta ataatacatg atgaattggg aagaggaact tctacctacg 2340
atggatttgg gttagcatgg gctatatcag aatacattgc acaaagatt ggtgcttttt 2400
gcatgtttgc aaccattttt catgaactta ctgccttggc caatcagata ccaactgtta 2460
ataatctaca tgtcacagca ctcaacctg aagagacctt aactatgctt tatcaggtga 2520
agaaaggtgt ctgtgatcaa agttttggga ttcattgttc agagcttgct aatttcccta 2580
agcatgtaat agagtgtgct aaacagaaag ccctggaact tgaggagttt cagtatatgt 2640
gagaatcgca aggatatgat atcatggaac cagcagcaa gaagtgtat ctggaaagag 2700
agcaagggtga aaaaattatt caggagtcc tgtccaagg gaaacaaatg ccctttactg 2760
aaatgtcaga agaaaacatc acaataaagt taaaacagct aaaagctgaa gtaaatgcaa 2820
agaataatag ctttgtaaat gaaatcattt cacgaataaa agttactacg tgaaaaatcc 2880
cagtaatgga atgaaggtaa tattgataag ctattgtctg taatagtttt atattgtttt 2940
atattaaccc tttttccata gtgttaactg tcagtgccca tgggtatca acttaataag 3000
atatttagta atattttact ttgaggacat tttcaaagat ttttattttg aaaaatgaga 3060

```

gctgtaactg aggactgttt gcaattgaca taggcaataa taagtgatgt gctgaatttt 3120  
 ataaataaaa tcatgtagtt tgtgg 3145

<210> 13

<211> 756

<212> PRT

<213> Homo sapiens

<400> 13

Met Ser Phe Val Ala Gly Val Ile Arg Arg Leu Asp Glu Thr Val Val  
 1 5 10 15

Asn Arg Ile Ala Ala Gly Glu Val Ile Gln Arg Pro Ala Asn Ala Ile  
 20 25 30

Lys Glu Met Ile Glu Asn Cys Leu Asp Ala Lys Ser Thr Ser Ile Gln  
 35 40 45

Val Ile Val Lys Glu Gly Gly Leu Lys Leu Ile Gln Ile Gln Asp Asn  
 50 55 60

Gly Thr Gly Ile Arg Lys Glu Asp Leu Asp Ile Val Cys Glu Arg Phe  
 65 70 75 80

Thr Thr Ser Lys Leu Gln Ser Phe Glu Asp Leu Ala Ser Ile Ser Thr  
 85 90 95

Tyr Gly Phe Arg Gly Glu Ala Leu Ala Ser Ile Ser His Val Ala His  
 100 105 110

Val Thr Ile Thr Thr Lys Thr Ala Asp Gly Lys Cys Ala Tyr Arg Ala  
 115 120 125

Ser Tyr Ser Asp Gly Lys Leu Lys Ala Pro Pro Lys Pro Cys Ala Gly  
 130 135 140

Asn Gln Gly Thr Gln Ile Thr Val Glu Asp Leu Phe Tyr Asn Ile Ala  
 145 150 155 160

Thr Arg Arg Lys Ala Leu Lys Asn Pro Ser Glu Glu Tyr Gly Lys Ile  
 165 170 175

Leu Glu Val Val Gly Arg Tyr Ser Val His Asn Ala Gly Ile Ser Phe  
 180 185 190

Ser Val Lys Lys Gln Gly Glu Thr Val Ala Asp Val Arg Thr Leu Pro  
 195 200 205

Asn Ala Ser Thr Val Asp Asn Ile Arg Ser Ile Phe Gly Asn Ala Val  
 210 215 220

Ser Arg Glu Leu Ile Glu Ile Gly Cys Glu Asp Lys Thr Leu Ala Phe  
 225 230 235 240

Lys Met Asn Gly Tyr Ile Ser Asn Ala Asn Tyr Ser Val Lys Lys Cys  
 245 250 255

Ile Phe Leu Leu Phe Ile Asn His Arg Leu Val Glu Ser Thr Ser Leu  
 260 265 270

Arg Lys Ala Ile Glu Thr Val Tyr Ala Ala Tyr Leu Pro Lys Asn Thr  
 275 280 285

His Pro Phe Leu Tyr Leu Ser Leu Glu Ile Ser Pro Gln Asn Val Asp  
 290 295 300

Val Asn Val His Pro Thr Lys His Glu Val His Phe Leu His Glu Glu  
 305 310 315 320

Ser Ile Leu Glu Arg Val Gln Gln His Ile Glu Ser Lys Leu Leu Gly  
 325 330 335

Ser Asn Ser Ser Arg Met Tyr Phe Thr Gln Thr Leu Leu Pro Gly Leu  
 340 345 350

Ala Gly Pro Ser Gly Glu Met Val Lys Ser Thr Thr Ser Leu Thr Ser  
 355 360 365

Ser Ser Thr Ser Gly Ser Ser Asp Lys Val Tyr Ala His Gln Met Val  
 370 375 380

Arg Thr Asp Ser Arg Glu Gln Lys Leu Asp Ala Phe Leu Gln Pro Leu  
 385 390 395 400

Ser Lys Pro Leu Ser Ser Gln Pro Gln Ala Ile Val Thr Glu Asp Lys  
 405 410 415

Thr Asp Ile Ser Ser Gly Arg Ala Arg Gln Gln Asp Glu Glu Met Leu  
 420 425 430

Glu Leu Pro Ala Pro Ala Glu Val Ala Ala Lys Asn Gln Ser Leu Glu  
 435 440 445

Gly Asp Thr Thr Lys Gly Thr Ser Glu Met Ser Glu Lys Arg Gly Pro  
 450 455 460

Thr	Ser	Ser	Asn	Pro	Arg	Lys	Arg	His	Arg	Glu	Asp	Ser	Asp	Val	Glu	465	470	475	480
Met	Val	Glu	Asp	Asp	Ser	Arg	Lys	Glu	Met	Thr	Ala	Ala	Cys	Thr	Pro	485	490	495	
Arg	Arg	Arg	Ile	Ile	Asn	Leu	Thr	Ser	Val	Leu	Ser	Leu	Gln	Glu	Glu	500	505	510	
Ile	Asn	Glu	Gln	Gly	His	Glu	Val	Leu	Arg	Glu	Met	Leu	His	Asn	His	515	520	525	
Ser	Phe	Val	Gly	Cys	Val	Asn	Pro	Gln	Trp	Ala	Leu	Ala	Gln	His	Gln	530	535	540	
Thr	Lys	Leu	Tyr	Leu	Leu	Asn	Thr	Thr	Lys	Leu	Ser	Glu	Glu	Leu	Phe	545	550	555	560
Tyr	Gln	Ile	Leu	Ile	Tyr	Asp	Phe	Ala	Asn	Phe	Gly	Val	Leu	Arg	Leu	565	570	575	
Ser	Glu	Pro	Ala	Pro	Leu	Phe	Asp	Leu	Ala	Met	Leu	Ala	Leu	Asp	Ser	580	585	590	
Pro	Glu	Ser	Gly	Trp	Thr	Glu	Glu	Asp	Gly	Pro	Lys	Glu	Gly	Leu	Ala	595	600	605	
Glu	Tyr	Ile	Val	Glu	Phe	Leu	Lys	Lys	Lys	Ala	Glu	Met	Leu	Ala	Asp	610	615	620	
Tyr	Phe	Ser	Leu	Glu	Ile	Asp	Glu	Glu	Gly	Asn	Leu	Ile	Gly	Leu	Pro	625	630	635	640
Leu	Leu	Ile	Asp	Asn	Tyr	Val	Pro	Pro	Leu	Glu	Gly	Leu	Pro	Ile	Phe	645	650	655	
Ile	Leu	Arg	Leu	Ala	Thr	Glu	Val	Asn	Trp	Asp	Glu	Glu	Lys	Glu	Cys	660	665	670	
Phe	Glu	Ser	Leu	Ser	Lys	Glu	Cys	Ala	Met	Phe	Tyr	Ser	Ile	Arg	Lys	675	680	685	
Gln	Tyr	Ile	Ser	Glu	Glu	Ser	Thr	Leu	Ser	Gly	Gln	Gln	Ser	Glu	Val	690	695	700	
Pro	Gly	Ser	Ile	Pro	Asn	Ser	Trp	Lys	Trp	Thr	Val	Glu	His	Ile	Val	705	710	715	720

Tyr Lys Ala Leu Arg Ser His Ile Leu Pro Pro Lys His Phe Thr Glu  
 725 730 735

Asp Gly Asn Ile Leu Gln Leu Ala Asn Leu Pro Asp Leu Tyr Lys Val  
 740 745 750

Phe Glu Arg Cys  
 755

<210> 14

<211> 2484

<212> DNA

<213> Homo sapiens

<400> 14

```

cttggtcttt ctggcgccaa aatgtcgttc gtggcagggg ttattcggcg gctggacgag 60
acagtgggtga accgcacgcg ggcgggggaa gttatccagc ggccagctaa tgctatcaaa 120
gagatgattg agaactgttt agatgcaaaa tccacaagta ttcaagtgat tgttaaagag 180
ggaggcctga agttgattca gatccaagac aatggcaccg ggatcaggaa agaagatctg 240
gatattgtat gtgaaagggt cactactagt aaactgcagt cctttgagga tttagccagt 300
attttctacct atggcttttc aggtgaggct ttggccagca taagccatgt ggctcatgtt 360
actattacaa cgaaaacagc tgatggaaag tgtgcataca gagcaagtta ctcatgtgga 420
aaactgaaag cccctcctaa accatgtgct ggcaatcaag ggaccacgat cacgggtggag 480
gacctttttt acaacatagc cagcaggaga aaagctttta aaaatccaag tgaagaatat 540
gggaaaattt tggaagttgt tggcagggtat tcagtaacaca atgcaggcat tagttttctca 600
gttaaaaaaac aaggagagac agtagctgat gttaggacac taccacatgc ctcaaccgtg 660
gacaatatct gctccatctt tggaaatgct gttagtcgag aactgataga aattggatgt 720
gaggataaaa ccctagcctt caaaatgaat ggttacatat ccaatgcaa ctactcagt 780
aagaagtgca tcttcttact ctcatcaac catcgtctgg tagaatcaac ttcttggaga 840
aaagccatag aaacagtgtg tgcagcctat ttgccccaaa acacacaccc attcctgtac 900
ctcagtttag aaatcagtc ccagaatgtg gatgttaatg tgcacccac aaagcatgaa 960
gttcacttcc tgcacgagga gagcatcctg gagcgggtgc agcagcacat cgagagcaag 1020
ctcctgggct ccaattcctc caggatgtac ttcaccacaga ctttgctacc aggacttgct 1080
ggccccctctg ggagatggt taaatccaca acaagtctga cctcgtcttc tacttctgga 1140
agtagtgata aggtctatgc ccaccagatg gttcgtacag attcccggga acagaagctt 1200
gatgcatttc tgcagcctct gagcaaacc ctgtccagtc agcccaggc cattgtcaca 1260
gaggataaga cagatatctt tagtggcagg gctaggcagc aagatgagga gatgcttgaa 1320
ctcccagccc ctgctgaagt ggctgccaaa aatcagagct tggaggggga tacaacaaag 1380
gggacttcag aaatgtcaga gaagagagga cctacttcca gcaaccccag aaagagacat 1440
cgggaagatt ctgatgtgga aatggtggaa gatgattccc gaaaggaaat gactgcagct 1500
tgtaccccc ggagaaggat cattaacctc actagtgttt tgagtctcca ggaagaaatt 1560
aatgagcagg gacatgaggt tctccgggag atgttgcata accactcctt cgtgggctgt 1620
gtgaatctc agtgggcctt ggcacagcat caaaccaagt tataccttct caacaccacc 1680
aagcttagtg aagaactgtt ctaccagata ctcatttatg attttgccaa ttttggtgtt 1740
ctcaggttat cggagccagc accgctcttt gaccttgcca tgcttgctt agatagtcca 1800
gagagtggct ggacagagga agatgggtccc aaagaaggac ttgctgaata cattgttgag 1860

```

```

tttctgaaga agaaggctga gatgcttgca gactatttct ctttggaat tgatgaggaa 1920
gggaacctga ttggattacc cttctgatt gacaactatg tgcccccttt ggagggactg 1980
cctatcttca ttcttcgact agccactgag gtgaattggg acgaagaaaa ggaatgtttt 2040
gaaagcctca gtaaagaatg cgctatgttc tattocatcc ggaagcagta catatctgag 2100
gagtcgaccc tctcaggcca gcagagtgaat gtgcctggct ccattccaaa ctccctggaag 2160
tggactgttg aacacattgt ctataaagcc ttgcgctcac acattctgcc tcctaaacat 2220
ttcacagaag atggaaatat cctgcagctt gctaacctgc ctgatctata caaagtcttt 2280
gagaggtgtt aaatatggtt atttatgcac tgtgggatgt gttcttcttt ctctgtattc 2340
cgatacaaag tgttgatca aagtgtgata tacaaagtgt accaacataa gtgttggtag 2400
cacttaagac ttatacttgc cttctgatag tattccttta tacacagtgg attgattata 2460
aataaataga tgtgtcttaa cata 2484

```

&lt;210&gt; 15

&lt;211&gt; 133

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 15

Met Glu Arg Ala Glu Ser Ser Ser Thr Glu Pro Ala Lys Ala Ile Lys

1

5

10

15

Pro Ile Asp Arg Lys Ser Val His Gln Ile Cys Ser Gly Gln Val Val

20

25

30

Leu Ser Leu Ser Thr Ala Val Lys Glu Leu Val Glu Asn Ser Leu Asp

35

40

45

Ala Gly Ala Thr Asn Ile Asp Leu Lys Leu Lys Asp Tyr Gly Val Asp

50

55

60

Leu Ile Glu Val Ser Asp Asn Gly Cys Gly Val Glu Glu Glu Asn Phe

65

70

75

80

Glu Gly Leu Thr Leu Lys His His Thr Ser Lys Ile Gln Glu Phe Ala

85

90

95

Asp Leu Thr Gln Val Glu Thr Phe Gly Phe Arg Gly Glu Ala Leu Ser

100

105

110

Ser Leu Cys Ala Leu Ser Asp Val Thr Ile Ser Thr Cys His Ala Ser

115

120

125

Ala Lys Val Gly Thr

130

&lt;210&gt; 16

<211> 426

<212> DNA

<213> Homo sapiens

<400> 16

```
cgaggcggat cgggtgttgc atccatggag cgagctgaga gctcgagtac agaacctgct 60
aaggccatca aacctattga tcggaagtca gtccatcaga tttgctctgg gcaggtggta 120
ctgagtctaa gcactgcggt aaaggagtta gtagaaaaca gtctggatgc tggtgccact 180
aatattgatc taaagcttaa ggactatgga gtggatctta ttgaagtttc agacaatgga 240
tgtggggtag aagaagaaaa cttcgaaggc ttaactctga aacatcacac atctaagatt 300
caagagtttg ccgacctaac tcaggttgaa acttttggct ttcgggggga agctctgagc 360
tcactttgtg cactgagcga tgtcaccatt tctacctgcc acgcacggc gaaggttgga 420
acttga                                           426
```



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/30588

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : Please See Extra Sheet.

US CL : 424/93.2, 130.1; 435 / 69.1, 410, 440; 536/24.5; 800/25

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/93.2, 130.1; 435 / 69.1, 410, 440; 536/24.5; 800/25

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN CAS, MEDLINE, BIOSIS, EMBASE, CAPLUS, WEST

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	KONG, Q. et al. PMS2-deficiency diminishes hypermutation of a lambda1 transgene in young but not older mice. Molecular Immunology. 1999, Vol. 36, pages 83-91, entire document.	1-4, 12-15, 22-25, 30-56
Y	VORA K.A. et al. Severe Attenuation of the B Cell Immune Response In Msh2-deficient Mice. Journal of Experimental Medicine. February 1999, Vol. 189, No. 3, pages 471-481, entire document.	1-2, 7-8, 12-13, 22-23, 30-56.
Y	WINTER, D.B. ET AL. Altered spectra of hypermutation in antibodies from mice deficient for the DNA mismatch repair protein PMS2. Proc. Natl. Acad. Sci., USA. June 1998, Vol. 95, pages 6953-6958, entire document.	1-4, 12-15, 22-25, 30-56

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

*	Special categories of cited documents:	"I"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E"	earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Z"	document member of the same patent family
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search	Date of mailing of the international search report
05 JANUARY 2001	12 MAR 2001

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-9230

Authorized officer  
David Saunders

Telephone No. (703) 308-0196

**INTERNATIONAL SEARCH REPORT**

International application No.

PCT/US00/30588

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SCHRADER, C.E. et al. Reduced Isotype Switching in Splenic B Cells From Mice Deficient in Mismatch Repair Enzymes. <i>Journal of Experimental Medicine</i> . 1999, Vol. 190, No. 3, pages 323-330, entire document.	1-4, 6, 12-13, 17 22-23, 27, 30-56

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/30588

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

A01N 63/00; A61K 39/395, 48/00; C07H 21/04; C12N 5/00, 15/00; C12P 21/06